- 1. Which of the following is mismatched pair?
 - A. Cellulose / structural polysaccharide in plant cells
 - B. Glycogen / Storage polysaccharide in animal cells
 - C. Amylose / Branched form of starch
 - D. Chitin / Structural polysaccharide contains nitrogen
- 2. In a polysaccharide (Glycosidic bond) is analog to (-----) in a polypeptide.
 - A. Hydrogen bond
 - B. Peptide bond
 - C. Phosphodiester
 - D. Ester bond
- 3. Each fat molecule:
 - A. Contains one glycerol and 3 fatty acids
 - B. Formed by dehydration reaction with removal of 3 water molecules.
 - C. Contains one glycerol and 2 fatty acids
 - D. Both A and B
 - E. Both A and C
- 4. Which levels of protein structure that least affected with disruption of hydrogen bonds?
 - A. Primary level
 - B. Secondary level
 - C. Tertiary level
 - D. Quaternary level
 - 5. Which of the following statements is correct?
 - A. Collagen is a globular protein made of 3 polypeptides
 - B. Hemoglobin is a globular protein made of 3 polypeptides
 - C. Collagen is a fibrous protein made of 4 polypeptides
 - D. Hemoglobin is globular protein made of 4 polypeptides

- 6. Describe the specific heat of water:
 - A. High
 - B. Low
 - C. Moderate
 - D. Equals 1 Cal (g.C)
 - E. Both A and D correct
- 7. How many water molecules needed to completely hydrolyze a polymer that is 11 monomers long?
 - A. 11
 - B. 10
 - C. 12
 - D. 9
 - E. Can't be determined
 - 8. In a polypeptide, the peptide bond formed between:
 - A. Carboxyl and amino group
 - B. Carboxyl and Carbonyl group
 - C. Carbonyl and amino group
 - D. Any of the above
 - E. None of the above
- 9. What type of covalent bond between amino acid side chains (R groups) functions in maintaining a polypeptide's specific three-dimensional shape?
 - A. ionic bond
 - B. hydrophobic interaction
 - C. van der Waals interaction
 - D. disulfide bond
 - E. hydrogen bond
- 10. If 14C-labeled uridine triphosphate is added to the growth medium of cells, what macromolecules will be labeled?
 - A. phospholipids
 - B. DNA
 - C. RNA
 - D. both DNA and RNA
 - E. proteins

- 11. Which of the following types of microscopes used to study internal ultrastructure of a cell?
 - A. SEM
 - B. TEM
 - C. Light microscopy
 - D. Phase- contrast microscopy
 - E. Super-resolution microscope
- 12. Which type of organelle or structure is primarily involved in the synthesis of oils, phospholipids, and steroids?
 - A. ribosome
 - B. lysosome
 - C. smooth endoplasmic reticulum
 - D. mitochondrion
 - E. contractile vacuole
 - 13. Which cell would be best for studying lysosomes?
 - A. muscle cell
 - B. leaf cell of a plant
 - C. nerve cell
 - D. phagocytic white blood cell
 - E. bacterial cell
 - 14. Which of the following factors would tend to increase membrane fluidity?
 - A. a greater proportion of unsaturated phospholipids
 - B. a lower temperature
 - C. a relatively high protein content in the membrane
 - D. a greater proportion of saturated phospholipids
 - E. a greater proportion of relatively large glycolipids compared with lipids having smaller molecular masses
 - 15. The major interaction responsible for stabilizing plasma membrane:
 - A. Hydrophilic interaction
 - B. Hydrophobic interaction
 - C. Ionic bonds
 - D. Hydrogen bonds

- 16. An animal cell lacking oligosaccharides on the external surface of its plasma membrane would likely be impaired in which function?
 - A. Transporting ions against an electrochemical gradient
 - B. Cell-cell recognition
 - C. Maintaining fluidity of the phospholipid bilayer
 - D. Attaching to the cytoskeleton
 - E. Establishing the diffusion barrier to charged molecules
 - 17. What are the membrane structures that function in active transport?
 - A. Peripheral proteins
 - B. Carbohydrates
 - C. Cholesterol
 - D. Cytoskeleton filaments
 - E. Integral proteins
- 18. What mechanisms do plants use to load sucrose produced by photosynthesis into specialized cells in the veins of leaves?
 - A. an electrogenic pump
 - B. a proton pump
 - C. cotransport protein
 - D. A and C only
 - E. A, B, and C
 - 19. Choose the correct word in each statement:

(DNA, RNA, Nucleotide, Nucleoside, Hydrogen bonds, Phosphodiester bonds)

- A. This bond stabilizes the double helix of DNA Hydrogen bonds
- B. This type of nucleic acid contains deoxyribose sugar and able to replicate itself DNA
- C. This bond found between the adjacent nucleotides within nucleic acid Phosphodiester
- D. Contains pentose sugar and nitrogenous base only Nucleoside
- E. This type of nucleic acid found as a single stranded molecule RNA

20.	About these typ	es of carb	ohydrates	, choose t	he correct	one in each	statement:
	(Chitin, Maltose,	Glucose.	Amylose.	Glycogen	. Cellulose	. Lactose. F	ructose)

- A. Structural polysaccharide contains B-glucose and cannot be digested by human Cellulose
- B. Disaccharide consists of 2 glucose monomers connected by 1,4 glycosidic linkage Maltose
- C. Monosaccharide represents the major fuel of cells Glucose
- D. Structural polysaccharide contains nitrogen Chitin
- E. Stored in liver and muscles Glycogen
- F. Ketose monosaccharide with formula (C₆H₁₂O₆) Fructose
- G. Milk sugar, that consists of glucose and galactose Lactose
- 21. The tendency of water molecules to stay close to each other as a result of hydrogen bonding _____.
- A. Provides the surface tension that allows leaves to float on water
- B. Is called cohesion
- C. Acts to moderate temperature
- D. Keeps water moving through the vessels in a tree trunk
- E. All of the listed responses are correct.
- 22. The amount of heat required to convert 1 g of any substance from the liquid to the gaseous state is defined as _____.
 - A. Molecular cohesion
 - B. 1 calorie
 - C. The specific heat of that substance
 - D. The heat of vaporization of that substance
 - E. The heat of fusion of that substance
- 23. Nonpolar molecules that cluster away from water molecules are called _____ molecules
 - A. Ionic
 - B. Saponified
 - C. Hydrophilic
 - D. Hydrophobic
 - E. None of the listed responses is correct.

- 24. Which of the following statements concerning unsaturated fats is true?
- A. They are more common in animals than in plants.
- B. They have double bonds in the carbon chains of their fatty acids.
- C. They generally solidify at room temperature.
- D. They contain more hydrogen than do saturated fats having the same number of carbon atoms.
- E. They have fewer fatty acid molecules per fat molecule.
- 25. Which class of biological polymer has the greatest functional variety?
- A. Polysaccharides
- B. Proteins
- C. DNA
- D. RNA
- 26. Among these biological polymers, which has the least structural variety?
- A. Polysaccharides
- B. Proteins
- C. DNA
- D. RNA
- 27. How will brief heating (to 95°C) affect macromolecular structures in aqueous solution?
 - A. DNA duplexes will unwind and separate.
 - B. Proteins will unfold (denature).
 - C. Starch will hydrolyze into monomeric sugars.
 - D. Proteins will hydrolyze into amino acids.
 - E. DNA duplexes will unwind and separate, and proteins will unfold (denature).
- 28. If cells are grown in a medium containing radioactive 15N, which of these molecules will be labeled?
 - A. Fatty acids only
 - B. Nucleic acids only
 - C. Proteins only
 - D. Amylase only
 - E. Both proteins and nucleic acids

- 29. If cells are grown in a medium containing radioactive 32P-labeled phosphate, which of these molecules will be labeled?
 - A. Phospholipids
 - B. Nucleic acids
 - C. Proteins
 - D. Amylose
 - E. Both phospholipids and nucleic acids
- 30. If a DNA sample were composed of 10% thymine, what would be the percentage of guanine?
- A. 10
- B. 20
- C. 40
- D. 80
- E. impossible to tell from the information given
- 31. Which of the following are nitrogenous bases of the pyrimidine type?
- A. guanine and adenine
- B. cytosine and uracil
- C. thymine and guanine
- D. ribose and deoxyribose
- E. adenine and thymine
- 32. Which of the following are nitrogenous bases of the purine type?
- A. cytosine and guanine
- B. guanine and adenine
- C. adenine and thymine
- D. thymine and uracil
- E. uracil and cytosine
- 33. Which of the following statements about the 5' end of a polynucleotide strand of DNA is correct?
- A. The 5' end has a hydroxyl group attached to the number 5 carbon of ribose.
- B. The 5' end has a phosphate group attached to the number 5 carbon of ribose.
- C. The 5' end has phosphate attached to the number 5 carbon of the nitrogenous base.
- D. The 5' end has a carboxyl group attached to the number 5 carbon of ribose.
- E. The 5' end is the fifth position on one of the nitrogenous bases.

- 34. What is the term used for a protein molecule that assists in the proper folding of other proteins?
 - A. Tertiary protein
 - B. Chaperonin
 - C. Enzyme protein
 - D. Renaturing protein
 - E. Denaturing protein
- 35. Misfolding of polypeptides is a serious problem in cells. Which of the following diseases are associated with an accumulation of misfolded polypeptides?
 - A. Alzheimer's only
 - B. Parkinson's only
 - C. Diabetes mellitus only
 - D. Alzheimer's and Parkinson's only
 - E. Alzheimer's, Parkinson's, and diabetes mellitus
- 36. At which level of protein structure are interactions between the side chains (R groups) most important?
 - A. Primary
 - B. Secondary
 - C. Tertiary
 - D. Quaternary
 - E. All of the above
 - 37. The tertiary structure of a protein is the
 - A. Bonding together of several polypeptide chains by weak bonds.
 - B. Order in which amino acids are joined in a polypeptide chain.
 - C. Unique three-dimensional shape of the fully folded polypeptide.
 - D. Organization of a polypeptide chain into an α helix or β pleated sheet.
 - E. Overall protein structure resulting from the aggregation of two or more polypeptide subunits.

- 38. Which of the following components that make a triglycerol molecule?
 - A. Alpha helix and beta pleated sheets
 - B. Glycogen and acetylglucose amine
 - C. Glycerol and fatty acids
 - D. Purine and ribose
 - E. Glucose and amino acids
- 39. Which is false for glucose and fructose
 - A. Both are ketoses
 - B. Both are monomers for sucrose
 - C. Both are structural isomers
 - D. Both are hexose
 - E. Both are monosaccharides
- 40. Smooth endoplasmic reticulum is responsible for:
 - A. Detoxifying drugs and poisons
 - B. Storage of calcium ions
 - C. Synthesizing sex hormones and lipids
 - D. None of the choices is correct
 - E. All choices are correct
- 41. Phosphodiester bonds are found in:
- A. Nucleoside
- B. Purine
- C. DNA strand
- D. Pyrimidine
- E. Fats
- 42. Water has maximum density at ----- C
- A. 4
- B. 0
- C. 37
- D. -4
- E. 100

- 43. The structural polysaccharide found in many insects and fungi is
- A. Cellulose
- B. Amylopectin
- C. Chitin
- D. Glycogen
- E. Amylose
- 44. What maintains the secondary structure of a protein?
- A. Peptide bonds
- B. Hydrogen bonds between the amino group of one peptide bond and the carboxyl group of another peptide bond
- C. Disulfide bonds
- D. Hydrophobic interactions
- E. Hydrogen bonds between the R groups
- 45. The reaction that break larger molecules into their smaller subunits is known as
- A. Polymerization reaction
- B. Dehydration reaction
- C. Condensation reaction
- D. Hydrolysis reaction
- E. None of the above
- 46. How many different kinds of polypeptides, each composed of 12 amino acids, could be synthesized using the 20 common amino acids?
 - A. 4¹²
 - B. 12²⁰
 - C. 240
 - D. 20
 - E. 20¹²
- 47. What type of carbohydrates does plant cells secret to hold (cement) the primary cell walls of adjacent cells together?
- A. Glycogen
- B. Pectin
- C. Amylose
- D. Amylopectin
- E. None of the above

- 48. Dehydration reactions are used in forming which of the following compounds?
- A. Triacylglycerides
- B. Polysaccharides
- C. Proteins
- D. Triacylglycerides and proteins only
- E. Triacylglycerides, polysaccharides, and proteins
- 49. Which of the following can pass through the pore complexes in the nuclear envelope?
 - A. Transport vesicles
 - B. Ribosomal proteins
 - C. RNA molecules
 - D. Ribosomal subunits
 - E. All choices are correct except transport vesicles
 - 50. The function of nucleolus is
 - A. Intracellular digestion
 - B. To manufacture polypeptides
 - C. To produce hydrogen peroxide
 - D. Store chromatin
 - E. To manufacture ribosomes
- 51. Which structure is the site of the synthesis of proteins that may be exported from the cell?
- A. Rough ER
- B. Plasmodesmata
- C. Golgi vesicles
- D. Lysosomes
- E. Free cytoplasmic ribosomes
- 52. Which plant cell organelle contain its own DNA and ribosomes?
- A. Glyoxysomes
- B. Peroxisomes
- C. vacuoles
- D. Chloroplasts

- 53. There are 20 different amino acids. What makes one amino acid different from another?
 - A. Different side chains (R groups) attached to a carboxyl carbon
 - B. Different side chains (R groups) attached to the amino groups
 - C. Different side chains (R groups) attached to an α carbon
 - D. Different structural and optical isomers
 - E. Different asymmetric carbons
- 54. What kind of chemical bond is found between paired bases of the DNA double helix?
 - A. Hydrogen
 - B. Ionic
 - C. Phosphodiester
 - D. Double or triple covalent bond
 - E. None of the above
 - 55. Pinocytosis is one type of:
 - A. Exocytosis
 - B. Endocytosis
 - C. Diffusion
 - D. Facilitated transport
 - E. Active transport
 - 56. Which organelle or structure is absent in plant cell:
 - A. Mitochondria
 - B. Golgi vesicles
 - C. Peroxisomes
 - D. Microtubules
 - E. Centrosomes
 - 57. Water molecules are able to form hydrogen bons with
 - A. Oils
 - B. Any compound that is not soluble in water
 - C. Oxygen gas
 - D. Chloride ions
 - E. Compounds that have polar covalent bonds

- 58. Water's high specific heat is mainly a consequence of the
- A. Absorption and release of heat when hydrogen bonds break and form
- B. Small size of the water molecules
- C. Fact that water is poor heat conductor
- D. Inability of water to dissipate heat into dry air
- E. High specific heat of oxygen and hydrogen atoms
- 59. Thylakoids, DNA and ribosomes are all components found in:
- A. Mitochondria
- B. vacuoles
- C. None of the options
- D. Chloroplasts
- E. Lysosomes
- 60. A cell with predominance of free ribosomes is most likely
- A. Producing primarily proteins for secretion
- B. Producing primarily cytoplasmic proteins
- C. Enlarging its vacuoles
- D. Digesting large food particles
- E. Constructing an extensive cell wall or ECM
- 61. The secretion of glycoproteins out of the cell is considered as an example of:
- A. Exocytosis
- B. Pinocytosis
- C. Phagocytosis
- D. Endocytosis
- E. Receptor mediated endocytosis
- 62. The molecules responsible for membrane transport are:
- A. Proteins
- B. Glycolipids
- C. Phospholipids
- D. Cholesterol
- E. Carbohydrate

- 63. Which of the following is a branched polysaccharide?
- A. Glycogen
- B. Amylose
- C. Cellulose
- D. Chitin
- E. None of the options
- 64. The concentration of calcium in a cell is 0.3%. The concentration of calcium in the surrounding fluid is 0.1%. How could the cell obtain more calcium?
 - A. Active transport
 - B. Pinocytosis
 - C. Osmosis
 - D. Simple diffusion
 - E. Facilitated diffusion
 - 65. Monomers made of pentose sugar, nitrogenous base and phosphate group are:
 - A. Fatty acids
 - B. Phospholipids
 - C. Amino acids
 - D. Nucleotides
 - E. Amylose
- 66. The substitution of glutamic acid with valine at 6th position of B-subunit of hemoglobin results in all of the following except:
 - A. Change in primary structure
 - B. Change in protein folding
 - C. Hemoglobin crystallization into a fiber
 - D. Increased efficiency of O2 transport by hemoglobin
 - E. Sickle cell disease in human
 - 67. All of the following nitrogenous bases found in DNA except:
 - A. Adenine
 - B. Uracil
 - C. Thymine
 - D. Cytosine

- D. Triacylglycerol
- E. Amylose
- 69. All of the following are a part of prokaryotic cell except:
- A. Ribosomes
- B. Nucleoid
- C. Cytoplasm
- D. Endoplasmic reticulum
- 70. Animal muscle cells adhere together strongly through ----- which are supported by intermediate filaments:
 - A. Desmosomes
 - B. Plasmodesmata
 - C. Tight junctions
 - D. Gap junctions
 - E. Cellulose fibers
 - 71. The monomers that make up amylopectin is:
 - A. Amino acid
 - B. Alpha glucose
 - C. Beta glucose
 - D. Fatty acid
 - E. Cellulose
 - 72. What kind of bonds hold water molecules together?
 - A. Hydrogen bonds
 - B. Ionic bonds
 - C. Hydrophilic bonds
 - D. Polar covalent bonds
 - E. None of the above

- 73. Which of the following organelles contains hydrolytic enzymes in animal cells?
- A. Glyoxysomes
- B. Central vacuole
- C. Peroxisomes
- D. Chloroplasts
- E. Lysosomes
- 74. Ions diffuse across membranes down their:
- A. Chemical gradients.
- B. Concentration gradients.
- C. Electrical gradients.
- D. Electrochemical gradients.
- E. A and B are correct
- 75. What mechanisms do plants use to load sucrose produced by photosynthesis into specialized cells in the veins of leaves?
 - A. n electrogenic pump
 - B. A proton pump
 - C. A cotransport protein
 - D. A and C only E
 - E. A, B, and C
- 76. Cell membranes are asymmetrical. Which of the following is a most likely explanation?
 - A. The cell membrane forms a border between one cell and another in tightly packed tissues such as epithelium.
 - B. Cell membranes communicate signals from one organism to another.
 - C. Cell membrane proteins are determined as the membrane is being packaged in the ER and Golgi.
 - D. The "innerness" and "outerness" of membrane surfaces are predetermined by genes.
 - E. Proteins can only span cell membranes if they are hydrophobic

- 77. Water passes quickly through cell membranes because
- A. The bilayer is hydrophilic.
- B. It moves through hydrophobic channels.
- C. Water movement is tied to ATP hydrolysis.
- D. it is a small, polar, charged molecule.
- E. it moves through aquaporins in the membrane
- 78. A cell whose cytoplasm has a concentration of 0.02 molar glucose is placed in a test tube of water containing 0.02 molar glucose. Assuming that glucose is not actively transported into the cell, which of the following terms describes the tonicity of the external solution relative to the cytoplasm of the cell?
 - A. Turgid
 - B. Hypertonic
 - C. Hypotonic
 - D. Flaccid
 - E. Isotonic
- 79. Which of the following statements correctly describes the normal tonicity conditions for typical plant and animal cells?
- A. The animal cell is in a hypotonic solution, and the plant cell is in an isotonic solution.
- B. The animal cell is in an isotonic solution, and the plant cell is in a hypertonic solution.
- C. The animal cell is in a hypertonic solution, and the plant cell is in an isotonic solution.
- D. The animal cell is in an isotonic solution, and the plant cell is in a hypotonic solution.
- E. The animal cell is in a hypertonic solution, and the plant cell is in a hypotonic solution
- 80. Which of the following membrane activities require energy from ATP hydrolysis?
- A. Facilitated diffusion.
- B. Movement of water into a cell
- C. Na+ ions moving out of the cell
- D. Movement of glucose molecules
- E. Movement of water into a paramecium

- 81. Which structure is not part of the endomembrane system?
- A. Nuclear envelope
- B. Chloroplast
- C. Golgi apparatus
- D. Plasma membrane
- E. ER
- 82. Which structure is common to plant and animal cells?
- A. Chloroplast
- B. Wall made of cellulose
- C. Central vacuole
- D. Mitochondrion
- E. Centriole
- 83. Cyanide binds with at least one molecule involved in producing ATP. If a cell is exposed to cyanide, most of the cyanide would be found within the
 - A. Mitochondria.
 - B. Ribosomes.
 - C. Peroxisomes.
 - D. Lysosomes.
 - E. Endoplasmic reticulum
- 84. Plasmodesmata in plant cells are most similar in function to which of the following structures in animal cells?
 - A. Peroxisomes
 - B. Desmosomes
 - C. Gap junctions
 - D. Extracellular matrix
 - E. Tight junctions

- 85. Ions can travel directly from the cytoplasm of one animal cell to the cytoplasm of an adjacent cell through
- A. Plasmodesmata
- B. intermediate filaments.
- C. Tight junctions.
- D. Desmosomes.
- E. Gap junctions
- 86. Which of the following are capable of converting light energy to chemical energy?
- A. Chloroplasts
- B. Mitochondria
- C. Leucoplasts
- D. Peroxisomes
- E. Golgi bodies
- 87. Organelles other than the nucleus that contain DNA include
- A. Ribosomes.
- B. Mitochondria.
- C. Chloroplasts.
- D. B and C only
- E. A, B, and C
- 88. Which of the following contains enzymes that transfer hydrogen from various substrates to oxygen?
 - A. lysosome
 - B. vacuole
 - C. mitochondrion
 - D. Golgi apparatus
 - E. peroxisome
 - 89. Grana, thylakoids, and stroma are all components found in
 - A. Vacuoles.
 - B. Chloroplasts.
 - C. Mitochondria.
 - D. lysosomes.

- 90. Which of the following contains hydrolytic enzymes?
- A. lysosome
- B. vacuole
- C. mitochondrion
- D. Golgi apparatus
- E. peroxisome
- 91. Which of the following is a compartment that often takes up much of the volume of a plant cell?
 - A. lysosome
 - B. vacuole
 - C. mitochondrion
 - D. Golgi apparatus
 - E. peroxisome
- 92. The liver is involved in detoxification of many poisons and drugs. Which of the following structures is primarily involved in this process and therefore abundant in liver cells?
 - A. rough ER
 - B. smooth ER
 - C. Golgi apparatus
 - D. Nuclear envelope
 - E. Transport vesicles
- 93. Which of the following produces and modifies polysaccharides that will be secreted?
- A. Lysosome
- B. Vacuole
- C. Mitochondrion
- D. Golgi apparatus
- E. Peroxisome

- 94. Tay-Sachs disease is a human genetic abnormality that results in cells accumulating and becoming clogged with very large and complex lipids. Which cellular organelle must be involved in this condition?
 - A. The endoplasmic reticulum
 - B. The Golgi apparatus
 - C. The lysosome
 - D. Mitochondria
 - E. Membrane-bound ribosomes
- 95. Which of the following correctly lists the order in which cellular components will be found in the pellet when homogenized cells are treated with increasingly rapid spins in a centrifuge?
- A. ribosomes, nucleus, mitochondria
- B. chloroplasts, ribosomes, vacuoles
- C. nucleus, ribosomes, chloroplasts
- D. vacuoles, ribosomes, nucleus
- E. nucleus, mitochondria, ribosomes
- 96. A primary objective of cell fractionation is to
- A. View the structure of cell membranes.
- B. identify the enzymes outside the organelles.
- C. determine the size of various organelles.
- D. separate the major organelles so that their particular functions can be determined.
- E. crack the cell wall so the cytoplasmic contents can be released.
- 97. Temperature is a measure of
- A. Specific heat.
- B. Average kinetic energy of molecules.
- C. Total kinetic energy of molecules.
- D. Celsius degrees.
- E. Joules.

98. Hydrophobic substances such as vegetable oil are

- A. nonpolar substances that repel water molecules.
- B. nonpolar substances that have an attraction for water molecules.
- C. polar substances that repel water molecules.
- D. polar substances that have an affinity for water.
- E. charged molecules that hydrogen-bond with water molecules
- 99. Which type of bond must be broken for water to vaporize?
 - A. ionic bonds
 - B. nonpolar covalent bonds
 - C. polar covalent bonds
 - D. hydrogen bonds
 - E. covalent bonds
- 100. Each water molecule is capable of forming
- A. one hydrogen bond.
- B. three hydrogen bonds.
- C. four hydrogen bonds.
- D. two covalent bonds and two hydrogen bonds.
- E. two ionic bonds and two hydrogen bonds.


1. Which of the following is true about this figure?

- A. It is a protein
- B. It is involved in the buffering of membrane fluidity
- C. Found only in plant cell membrane
- D. Can be used to make other molecules such as sex hormones
- E. Both B and C correct

2. Which of the following true about this figure?

- A. It represents nucleoside
- B. Called nucleotide or nucleoside monophosphate
- C. Can found in both DNA or RNA
- D. Both A and B
- E. Both B and C


3. According to this figure:



- A. (A) represents: Food vacuole, lysosome, damaged organelle
- B. (B) represents: Food vacuole, lysosome, damaged organelle
- C. The overall process called: Autophagy, phagocytosis, exocytosis
- D. If the enzymes of this organelle are missing or defective, the result could be: Lysosomal storage disease such as cystic fibrosis Lysosomal storage disease such as Tay-sachs disease Alzheimer disease

Parkinson disease

- E. The best PH for the enzymes that found in these organelles: Acidic, basic, neutral
- F. Are these organelles part of endomembrane system? Yes, NO
- 4. This figure shows four levels of protein structure, choose the correct one in each statement:



- A. Which level represents linear chain of amino acids joined by peptide bond? 1
- B. Which level is the least affected by disruption of hydrogen bonds? 1
- C. Which level represents regions stabilized by hydrogen bonds between backbone? 2
- D. Which level represents aggregation of two or more polypeptides? 4
- E. Which level involve the formation of disulfide bridge between two cystine monomers? 3

Plasma membrane
Pseudopodium
Phagosome (food vacuole)

B

Vesicle
Coated pit
Receptor
Coated pit
Receptor
Coated vesicle


- 5. The figure shows 3 types of endocytosis, choose the correct type in each statement:
 - A. Represent engulfing of particles and formation of food vacuole A
 - B. LDL can enter the cells by this type C
 - C. Represents non-specific endocytosis in which droplets of dissolved material enter the cell B

6. Which of the following true about this figure?




- A. It results from hydrogen bonding between water molecules
- B. It called specific heat
- C. It called surface tension
- D. It results from hydrogen bonding between the water at interface and the air above
- E. Both A and C
- F. Both B and D

7. Which of the following is true about this figure




- A. It is alpha glucose that contains nitrogen and found in cellulose
- B. It is beta glucose that contains nitrogen and found in cellulose
- C. It is alpha glucose that contains nitrogen and found in chitin
- D. It is beta glucose that contains nitrogen and found in chitin


8. According to this figure:



- In order to make one of theses molecules we need:
 - A. Glycerol 2 fatty acids and choline
 - B. Glycerol 2 fatty acids and phosphate
 - C. Addition of 3 water molecules
 - D. Glycerol 3 fatty acids with removal of 3 water molecules
 - E. None of the above
- Which of them can be found solid at room temperature? B
- Plants fats and oils are examples of which type? A

9. Which of the following correct about these organelles?






- A. Both are part of endomembrane system
- B. Both can convert solar energy to chemical energy
- C. Both surrounded by two membranes separated by intermembrane space
- D. Both contains its own DNA, ribosomes and enzyme
- E. Only C and D
- F. Only A and B

10. Which of the following not part of extracellular matrix (ECM)?

- A. Collagen and integrin
- B. Cellulose and pectin
- C. Integrin and fibronectin
- D. Collagen and proteoglycan




11. Which of the following is true about middle lamella?

A. It made of pectin to connect adjacent primary cell walls of plant cells


- B. It made of cellulose to connect adjacent primary cell walls of plant cells
- C. Its function to maintain the shape of nucleus
- D. None of the above

12. According to this figure:



- A. Which bond represents polar covalent bond? A or B
- B. Which bond represents hydrogen bonds? A or B
- C. Each water molecule can form 4 hydrogen bonds with other molecules.
- D. To vaporize water, which bond must be broken? A or B
- 13. Which of the following is a common feature between all cells?
- A. Nucleus
- B. Chromosome
- C. Cell membrane
- D. Cytosol
- E. All of the except of A
- 14. Which of the following is true about nuclear lamina?
- A. It made of intermediate filament
- B. Its function to maintain the shape of nucleus
- C. It is a framework of protein fibers extending through the nucleus
- D. Both A and B
- E. Both B and C


15. According to this figure



- Which of them is a fibrous protein? A
- Which of them is a globular protein? B


 If there is a change in the primary structure of (B), This will cause a disease known as Sickle-cell anemia

16. According to this figure, choose the correct junction between these:




- A. Which of the following prevents leakage of extracellular fluid across a layer of cells? A
- B. Which of the following provides cytoplasmic channels from one cell to another? C
- C. Plasmodesmata in plant cells is similar to which structure in this figure? C
- D. Which of the following anchored to cytoplasm by intermediate filament? B
- 17. Which of the following not function of smooth endoplasmic reticulum?
- A. Storage of calcium ions
- B. Detoxification of drugs and poisons
- C. Making proteins that will be secreted out of the cell
- D. Metabolism of carbohydrate

18. According to this figure:



- A. (A) represents proton pump
- B. (B) represent H+/sucrose cotransporter
- C. Name these processes Cotransport
- D. Major electrogenic pump in animal cells is Na+/K+ pump and in plant cells is proton pump
- E. Why it called electrogenic? Because it contributes in membrane potential
- F. Voltage across membrane called Membrane potential
- G. We need ATP as a source of energy in (Active or passive transport)
- H. Water moves quickly across membrane due to Aquaporins
- CO2, O2 and other nonpolar molecules can pass through membrane by simple diffusion
- 19. About sodium-potassium pump decide whether these statements true or false?
 - A. It is an active process requires ATP as a source of energy (T)
 - B. Pumps 3 sodium ions into the cell (F)
 - C. Pumps 2 potassium ions into the cell (T)
 - D. It is major electrogenic pump in plant cells (F)
- 20. Ions diffuse across membrane down there:
- A. Chemical gradient
- B. Electrical gradient
- C. Electrochemical gradient
- D. None of the above

21. The figure shows animal cells placed in 3 solutions, choose the correct one:



- A. Which letter represents the normal tonicity of animal cell? A
- B. In (C), cell will lose water so it will ----- (Shrink, lysed, normal)
- C. The healthy state of plant cell in ----- (Hypotonic / Hypertonic/ Isotonic)
- D. What we mean be osmosis? Diffusion of water molecules form region of high free water concentration to region of low free water concentration
- 22. Write the molecular formula of a polymer contains 10 ribose molecules.

- 23. Solution in which water is the solvent are called
- A. non aqueous solution
- B. aqueous solution
- C. water solution
- D. None of the above
- 24. Insoluble fibers refer to:
- A. Cellulose
- B. Chitin
- C. Glycogen
- D. Amylose
- E. None of the above

25. Which of the following mismatched pair?

- A. Nucleolus / Production of ribosomes
- B. Mitochondria / ATP production
- C. Chloroplasts / Photosynthesis
- D. Microtubules / Formation of nuclear lamina
- E. Lysosome / Intracellular digestion

Table 6.1 The Structure and Function of the Cytoskeleton


Property	Microtubules (Tubulin Polymers)	Microfilaments (Actin Filaments)	Intermediate Filaments
Structure	Hollow tubes	Two intertwined strands of actin	Fibrous proteins coiled into cables
Diameter	25 nm with 15-nm lumen	7 nm	8–12 nm
Protein subunits	Tubulin, a dimer consisting of α -tubulin and β -tubulin	Actin	One of several different proteins (such as keratins)
Main functions	Maintenance of cell shape (compression-resisting "girders"); cell motility (as in cilia or flagella); chromosome movements in cell division; organelle movements	Maintenance of cell shape (tension- bearing elements); changes in cell shape; muscle contraction; cytoplasmic streaming in plant cells; cell motility (as in amoeboid movement); division of animal cells	Maintenance of cell shape (tension- bearing elements); anchorage of nucleus and certain other organ- elles; formation of nuclear lamina
Fluorescence micro- graphs of fibroblasts. Fibroblasts are a favorite cell type for cell biology studies. In each, the structure of interest has been tagged with fluorescent molecules. The DNA in the nucleus has also been tagged in the first micrograph (blue) and third micrograph (orange).	Column of tubulin dimers	Actin subunit	Keratin proteins Fibrous subunit (keratins coiled together)

26. Which of the following does not contains amino acids?


- A. Insulin
- B. Antibodies
- C. Cholesterol
- D. Hemoglobin

- 27. About proteins, decide if these statements are correct:
- A. In a polypeptide, amino acids arranged in a branched polymer F
- B. We can find hydrophobic amino acids at the surface of a globular protein F
- C. Disulfide bridges is a covalent bond responsible of stabilization of secondary level of protein F
- D. When the protein loses its native shape, the process called renaturation F
- 28. Which of the following found only in RNA?
- A. Guanine and ribose
- B. Adenine and deoxyribose
- C. Uracil and deoxyribose
- D. Uracil and ribose
- E. None of the above

29. Which of the following is false about this figure?



- A. It is a prokaryotic cell with membrane bound nucleus
- B. It is a bacterium
- C. It is a unicellular organism
- D. It DNA located with non-enclosed region called nucleoid
- E. None of the above




30. About cell membrane (Plasma membrane), which of the following false statement?


- A. The components of cell membrane held together by hydrophobic interaction
- B. Phospholipids can move rapidly by lateral movements
- C. Most proteins are held in place by attachment to cytoskeleton and ECM
- D. There is a high level of cholesterol in cell membrane of plant cells
- E. Phospholipids represents the main fabric of membrane, while protein determines its function

31. Which of the following acts as Fluidity buffer within the membrane?


- A. Cholesterol
- B. Phospholipids
- C. Proteins
- D. Carbohydrate

Unsaturated versus saturated hydrocarbon tails





Cholesterol within the animal cell membrane



Cholesterol reduces membrane fluidity at moderate temperatures, but at low temperatures hinders solidification.

- 32. Which of the following molecules involved in cell-cell regonition?
- A. Cholesterol
- B. Carbohydrate
- C. Phospholipids
- D. Proteins
- E. None of the above
- 33. About DNA, which of the following is fasle:
- A. DNA is a double helix
- B. Adenine forms a complemetery pairing with thymine by 2 hydrogen bonds
- C. Guanine forms a complemetery pairing with cytosine by 3 hydrogen bonds
- D. 3' end contains phosphate wihle 3' ends contains OH
- E. DNA can replictaes itself

- 34. Which of the following transporter proteins involved in active transport:
- A. Only channel protein
- B. Only carrier protein
- C. Both of carrier and channel
- D. There is no need of transporter protein in active tranport
- E. None of the above
- 35. Which of the following is a measure of clarity:
- A. Magnification
- B. Contrast
- C. Resolution
- D. Both of reolution and contrast
- E. Both of magnification of contrast
- 36. About microscopes, choose the false statement:
- A. We use TEM to study internal structure of cells
- B. We use SEM to get 3D image
- C. LM uses visible light
- D. In EM we ues glass lenses
- E. We can study living cells with LM
- 37. The volume enclosed by the plasma membrane of plant cells is often much larger than the corresponding volume in animal cells. The most reasonable explanation for this observation is that
- A. plant cells are capable of having a much higher surface-to-volume ratio than animal cells.
- B. plant cells have a much more highly convoluted (folded) plasma membrane than animal cells.
- C. plant cells contain a large vacuole that reduces the volume of the cytoplasm.
- D. animal cells are more spherical, while plant cells are elongated.
- E. the basic functions of plant cells are very different from those of animal cells

- 38. Which of the following is a compartment that often takes up much of the volume of a plant cell?
- A. lysosome
- B. vacuole
- C. mitochondrion
- D. Golgi apparatus
- E. Peroxisome
- 39. The Golgi apparatus has a polarity or sidedness to its structure and function. Which of the following statements correctly describes this polarity?
- A. Transport vesicles fuse with one side of the Golgi and leave from the opposite side.
- B. Proteins in the membrane of the Golgi may be sorted and modified as they move from one side of the Golgi to the other.
- C. Lipids in the membrane of the Golgi may be sorted and modified as they move from one side of the Golgi to the other.
- D. Soluble proteins in the cisternae (interior) of the Golgi may be sorted and modified as they move from one side of the Golgi to the other.
- E. All of the above correctly describe polar characteristics of the Golgi function.
- 40. Organelles other than the nucleus that contain DNA include
 - A. Ribosomes.
 - B. mitochondria.
 - C. chloroplasts.
 - D. B and C only
 - E. A, B, and C
- 41. Why isn't the mitochondrion classified as part of the endomembrane system?
- A. It only has two membrane layers.
- B. Its structure is not derived from the ER.
- C. It has too many vesicles.
- D. It is not involved in protein synthesis.
- E. It is not attached to the outer nuclear envelope

- 42. Which structure is common to plant and animal cells?
 - A. chloroplast
 - B. wall made of cellulose
 - C. central vacuole
 - D. mitochondrion
 - E. centriole
- 43. Which of the following is present in a prokaryotic cell?
- A. mitochondrion
- B. ribosome
- C. nuclear envelope
- D. chloroplast
- E. ER
- 44. In what way do the membranes of a eukaryotic cell vary?
- A. Phospholipids are found only in certain membranes.
- B. Certain proteins are unique to each membrane.
- C. Only certain membranes of the cell are selectively permeable.
- D. Only certain membranes are constructed from amphipathic molecules.
- E. Some membranes have hydrophobic surfaces exposed to the cytoplasm, while others have hydrophilic surfaces facing the cytoplasm.