



Lipids

Summer 2023-2024

Lipids




- Lipids are a heterogeneous class of naturally occurring organic compounds that share some properties based on structural similarities, mainly a dominance of nonpolar groups.
- They are Amphipathic.
- They are insoluble in water, but soluble in fat or organic solvents (ether, chloroform, benzene, acetone).
- They are widely distributed in plants & animals.

Classes



- Simple lipids (fats, oils, and waxes)
- Complex lipids (glycerides , glycerophospholipids, sphingolipids, glycolipids, lipoproteins)
- Derived lipids (fatty acids, alcohols, eicosanoids)
- Cyclic lipids (steroids)



Lipid Functions



- Lipids include:
 - **Storage lipids for energy purpose**
 - They are storable to unlimited amounts (vs. carbohydrates)
 - They provide a considerable amount of energy to the body (25% of body needs) & provide a high-energy value (more energy per gram vs. carbohydrates & proteins)
 - **Structural lipids in membranes**
- Signaling molecules, hormone precursors, cofactors, & pigments
- Shock absorbers and thermal insulators




Fatty acids



- Aliphatic mono-carboxylic acids
- Formula: $R-(CH_2)_n-COOH$
- Lengths
 - Physiological (12-24)
 - Abundant (16 and 18)
- Degree of unsaturation
- Amphipathic molecules


Functions:

- Building blocks of other lipids
- Modification of many proteins (lipoproteins)
- Important fuel molecules
- Derivatives of important cellular molecules




Types of fatty acids


- Saturated fatty acids are those with all of the C-C bonds being single.
- Unsaturated fatty acids are those with one or more double bonds between carbons:
 - Monounsaturated fatty acid: a fatty acid containing one double bond.
 - Polyunsaturated fatty acids contain two or more double bonds.





Cis vs. trans bonds



linoleic acid: *trans* configuration (*trans* isomer)



linoleic acid: *cis* configuration (*cis* isomer)





Physiologically:

- cis isomer predominates
- trans is rare

Properties of fatty acids



- The properties of fatty acids (melting point) are dependent on chain length and degree of saturation.



Properties of saturated fatty acids



Short chain F.A. (2-4)	Medium-chain F.A. (6-10)	Long chain F.A. (12-20)
They are liquid in nature	Solids at room temperature	Solids at room temperature
Water-soluble	Water-soluble	Water-insoluble
Volatile at RT	Non-volatile at RT	Non-volatile
Acetic, butyric, caproic acids	Caprylic & capric acids	Palmitic and stearic acids



Greek number prefix




Number	prefix	Number	prefix	Number	prefix
1	Mono-	5	Penta-	9	Nona-
2	Di-	6	Hexa-	10	Deca-
3	Tri-	7	Hepta-	20	Eico-
4	Tetra-	8	Octa-		


Naming of a fatty acid



- Alkane to oic
 - Octadecane (octa and deca) is octadecanoic acid**
 - One double bond = octadecenoic acid
 - Two double bonds = octadecadienoic acid
 - Three double bonds = octadecatrienoic acid
- Designation of carbons and bonds
 - 18:0 = a C18 fatty acid with no double bonds**
 - stearic acid (18:0); palmitic acid (16:0)
 - 18:2 = two double bonds (linoleic acid)**
- Designation of the location of bonds
 - Δn : The position of a double bond**
 - cis- $\Delta 9$: a cis double bond between C 9 and 10
 - trans- $\Delta 2$: a trans double bond between C 2 and 3




Palmitoleic acid ($\omega 7, 16:1, \Delta^9$)




Oleic acid ($\omega 9, 18:1, \Delta^9$)




*Linoleic acid ($\omega 6, 18:2, \Delta^{9,12}$)



* α -Linolenic acid ($\omega 3, 18:3, \Delta^{9,12,15}$)



*Arachidonic acid ($\omega 6, 20:4, \Delta^{5,8,11,14}$)



Eicosapentaenoic acid ($\omega 3, 20:5, \Delta^{5,8,11,14,17}$)




Number of carbons	Number of double bonds	Common name	Systematic name	Formula
14	0	Myristate	n-Tetradecanoate	$\text{CH}_3(\text{CH}_2)_{12}\text{COO}^-$
16	0	Palmitate	n-Hexadecanoate	$\text{CH}_3(\text{CH}_2)_{14}\text{COO}^-$
18	0	Stearate	n-Octadecanoate	$\text{CH}_3(\text{CH}_2)_{16}\text{COO}^-$
18	1	Oleate	cis- Δ^9 -Octadecenoate	$\text{CH}_3(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_7\text{COO}^-$
18	2	Linoleate	cis,cis- Δ^9,Δ^{12} -Octadecadienoate	$\text{CH}_3(\text{CH}_2)_4\text{CH}=\text{CHCH}_2\text{CH}(\text{CH}_2)_7\text{COO}^-$
18	3	Linolenate	all-cis- $\Delta^9,\Delta^{12},\Delta^{15}$ -Octadecatrienoate	$\text{CH}_3\text{CH}_2(\text{CH}=\text{CHCH}_2)_3(\text{CH}_2)_6\text{COO}^-$
20	4	Arachidonate	all-cis- $\Delta^5,\Delta^8,\Delta^{11},\Delta^{14}$ -Eicosatetraenoate	$\text{CH}_3(\text{CH}_2)_4(\text{CH}=\text{CHCH}_2)_4(\text{CH}_2)_2\text{COO}^-$

Another way of naming






- (ω)-C: distal methyl C as #1





Per softgel		
Organic Flaxseed Oil (Linseed Oil)		400mg
Pure Fish Oil		400mg
Starflower Oil (Borage Oil)		400mg
typically providing:		
Alpha Linolenic Acid (ALA)	Omega-3	200mg
Docosahexaenoic Acid (DHA)	Omega-3	48mg
Eicosapentaenoic Acid (EPA)	Omega-3	72mg
Gamma Linolenic Acid (GLA)	Omega-6	88mg
Linoleic Acid (LA)	Omega-6	204mg
Oleic Acid	Omega-9	168mg

Omega-3 fatty acids









Omega-6 fatty acids



- **Essential FAs:**
 - Linoleic acid: precursor of arachidonates
 - Linolenic acid: precursor of EPA and DHA




Numerical Symbol	Common Name and Structure	Comments
18:1 ^{Δ9}	Oleic acid 	Omega-9 monounsaturated
18:2 ^{Δ9,12}	Linoleic acid 	Omega-6 polyunsaturated
18:3 ^{Δ9,12,15}	α-Linolenic acid (ALA) 	Omega-3 polyunsaturated
20:4 ^{Δ5,8,11,14}	Arachidonic acid 	Omega-6 polyunsaturated
20:5 ^{Δ5,8,11,14,17}	Eicosapentaenoic acid (EPA) 	Omega-3 polyunsaturated (fish oils)
22:6 ^{Δ4,7,10,13,16,19}	Docosahexaenoic acid (DHA) 	Omega-3 polyunsaturated (fish oils)

Derived fatty acids: Eicosanoids


Arachidonate



all *cis*- $\Delta^5, \Delta^8, \Delta^{11}, \Delta^{14}$ -eicosatetraenoate, $\text{CH}_3(\text{CH}_2)_4(\text{CH}=\text{CHCH}_2)_4(\text{CH}_2)_2\text{COO}^-$



Arachidonic acid






Eicosanoids and their functions




They control cellular function in response to injury

- Prostaglandins
 - Induction of inflammation
 - Inhibition of platelet aggregation
 - Inhibition of blood clotting
- Leukotrienes
 - Constriction of smooth muscles
 - Asthma
- Thromboxanes
 - Constriction of smooth muscles
 - Induction of platelet aggregation
- Prostacyclins
 - An inhibitor of platelet aggregation
 - Induction of vasodilation

Neither memorize the specific functions nor the structures



Aspirin is good




COX: Cyclooxygenase

Targets of Aspirin



- Cyclooxygenase is present in three forms in cells, COX-1, COX-2, and COX-3.
- Aspirin targets both, but COX-2 should only be the target.




Aspirin can be bad

Cardiovascular disease vs. bleeding



WARNING

- Aspirin also causes excessive bleeding among the elderly.




CAUTION

Age-specific risks, severity, time course, and outcome of bleeding on long-term antiplatelet treatment after vascular events: a population-based cohort study

Linxin Li*, Olivia C Geraghty*, Ziyah Mehta, Peter M Rothwell, on behalf of the Oxford Vascular Study

Interpretation In patients receiving aspirin-based antiplatelet treatment without routine PPI use, the long-term risk of major bleeding is higher and more sustained in older patients in practice than in the younger patients in previous trials, with a substantial risk of disabling or fatal upper gastrointestinal bleeding. Given that half of the major bleeds in patients aged 75 years or older were upper gastrointestinal, the estimated NNT for routine PPI use to prevent such bleeds is low, and co-prescription should be encouraged.



- A new generation drug, Celebrex, targets COX2, but is prescribed with a strong warning of side effects on the label.



Cardiovascular Risk

- CELEBREX may cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. All NSAIDs may have a similar risk. This risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk. (See **WARNINGS** and **CLINICAL TRIALS**).
- CELEBREX is contraindicated for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery (see **WARNINGS**).

Omega fatty acids





- Omega-3 fatty acids
 - α -linolenic acid $\rightarrow$ eicosapentaenoic acid (EPA) $\rightarrow$ docosahexaenoic acid (DHA)
 - They reduce inflammatory reactions.
- Omega-6 fatty acids:
 - Arachidonic acid
- Omega-9 fatty acids
 - Oleic acid
 - It reduces cholesterol in the circulation.


**Healthy
mediterranean
food**




Complex lipids




Triglycerides



Types of glycerides



Tristearin
a simple triglyceride




a mixed triglyceride

How soluble will a triglyceride be if fatty acids are unsaturated?


Solid vs. liquid fats



- Vegetable oils consist almost entirely of unsaturated fatty acids, whereas animal fats contain a much larger percentage of saturated fatty acids.
 - This is the primary reason for the different melting points of fats and oils.



A fat




An oil

Saponification




- Hydrolysis: steam, acid, enzyme (e.g., lipase of pancreas)
- Saponification: Alkaline hydrolysis produces salts of fatty acids (soaps). Soaps cause emulsification of oily material.



How does soap work?




- When mixed with water, the hydrophobic hydrocarbon tails cluster together to create a nonpolar microenvironment and the hydrophilic ionic heads interact with water.
- The resulting spherical clusters are called micelles.
- Grease and dirt are trapped inside micelles and the complex can be rinsed away.



Hydrogenation



- The carbon-carbon double bonds in vegetable oils can be hydrogenated to yield saturated fats in the same way that any alkene can react with hydrogen to yield an alkane.




Trans fat



- Although the animal fat is unhealthy, it has better cooking properties and better taste.
- Therefore, chemists invented a method of converting unsaturated oil into solid form by partially hydrogenating it.
- Partial hydrogenation converts some, but not all, double bonds into single bonds generating (trans fats).

The primary health risk identified for trans fat consumption is an elevated risk of coronary heart disease (CHD).



Example: margarine




- In margarine, only about two-thirds of the double bonds present in the starting vegetable oil are hydrogenated, so that the margarine remains soft in the refrigerator and melts on warm toast.

Nutrition Facts		
Serving Size 1 Tbsp (14g)		
Servings Per Container 32		
Amount Per Serving		
Calories	100	Calories from Fat 100
%		
		% Daily Value*
Total Fat	11g	17%
Saturated Fat	2g ←	10%
Trans Fat	3g ←	
Cholesterol	0mg	→ 0%



Waxes



- Solid simple lipids containing a monohydric alcohol (C16 ~ C30, higher molecular weight than glycerol) esterified to long-chain fatty acids (C14 ~ C36).
 - **Example: palmitoyl alcohol**
- Insoluble in water
- Not easily hydrolyzed (fats) & indigestible
- Very resistant to rancidity
- No nutritional value
- External coating of plant leaves that prevent loss of water




Type	Structural Formula	Source	Uses
Beeswax	$\text{CH}_3(\text{CH}_2)_{14}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-(\text{CH}_2)_{29}\text{CH}_3$	Honeycomb	Candles, shoe polish, wax paper
Carnauba wax	$\text{CH}_3(\text{CH}_2)_{24}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-(\text{CH}_2)_{29}\text{CH}_3$	Brazilian palm tree	Waxes for furniture, cars, floors, shoes
Jojoba wax	$\text{CH}_3(\text{CH}_2)_{18}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-(\text{CH}_2)_{19}\text{CH}_3$	Jojoba	Candles, soaps, cosmetics

Do not memorize the structures but study them

Membrane lipids




The most prevalent class of lipids in membranes is the glycerophospholipids



Phospholipids (phosphoacylglycerols)



- Phosphatidic acids
- Phosphatidylcholine (lecithin)
 - Most abundant membrane lipid
- Cephalins
 - Phosphatidylethanolamine
 - Phosphatidylserine
 - abundant in brain
- Phosphatidylinositol
 - sends messages across cell membranes
- Cardiolipin
- Plasmalogens



Phosphatidic acid	—	— H
Phosphatidylethanolamine	Ethanolamine	— CH ₂ —CH ₂ —NH ₃ ⁺
Phosphatidylcholine	Choline	— CH ₂ —CH ₂ —N ⁺ (CH ₃) ₃
Phosphatidylserine	Serine	— CH ₂ —CH—NH ₃ ⁺ COO ⁻

Glycerophospholipids - Lecithins



- Snake venom contain lecithinase, which hydrolyzes polyunsaturated fatty acids and converting lecithin into lysolecithin
 - hemolysis of RBCs

Phosphatidylcholine


Choline



Emulsification



- Because of their amphipathic nature, they act as emulsifying agents, that is substances that can surround nonpolar molecules and keep them in suspension in water.




Glycerophospholipids - Cardiolipins



- Diphosphatidyl-glycerol
- Found in the inner membrane of mitochondria
- Initially isolated from heart muscle (cardio)
- Structure: 3 molecules of glycerol, 4 fatty acids & 2 phosphate groups



**Do not memorize
the structures but
study them**



Plasmalogens




- They are found in the cell membrane phospholipids fraction of brain & muscle, liver, and semen.
- They have a protective role against reactive oxygen species
- Structure:
 - Precursor: Dihydroxyacetone phosphate
 - Unsaturated fatty alcohol at C1 connected by ether bond
 - In mammals: at C3; phosphate + ethanolamine or choline
- Major classes of plasmalogens
 - Ethanolamine plasmalogen (myelin-nervous tissues)
 - Choline plasmalogen (cardiac tissue)
 - Platelet activating factor
 - Serine plasmalogens




Glycerophospholipids - Inositides




- Phosphatidyl inositol
- Nitrogenous base: cyclic sugar alcohol (inositol)
- Structure: glycerol, saturated FA, unsaturated FA, phosphoric acid, & inositol
- Source: Brain tissues
- Functions:
 - Major component of cell membrane
 - Signalling molecules are produced upon hydrolysis



The different structures of phospholipids




Uses of liposomes: delivery



Sphingolipids



- Sphingolipids are found in the plasma membranes of all eukaryotic cells and is highest in the cells of the central nervous system
- The core of sphingolipids is the long-chain amino alcohol, sphingosine




Mysterious lipids




Sphynx → sphingolipids

Trivia

Named for the Sphinx of Thebes, who killed passersbies that could not solve her riddles




Ceramide





Types of sphingolipids



- The sphingolipids are divided into two subcategories:
 - **Sphingomyelin**
 - It is a sphingolipid that is a major component of the coating around nerve fibers.
 - The group attached to C1 is a phosphocholine
 - **Glycosphingolipid (or glycolipids)**




Zooming into the myelin



Glycolipids




- Sphingolipids can also contain carbohydrates attached at C-1 and these are known as glycolipids
- Glycolipids are present on cell membranes and act as cell surface receptors that can function in cell recognition (e.g., pathogens) and chemical messengers
- There are three types of glycolipids
 - Cerebrosides
 - Globosides
 - Gangliosides




Glycolipids



- **Cerebrosides:** the simplest glycolipids, contain a single hexose (galactose or glucose).
- **Globosides and gangliosides** are more complex glycolipids.
- Both contain glucose, galactose, and N-acetylgalactosamine, but gangliosides must also contain sialic acid.





Gangliosides are bound by cholera toxin in the human intestine facilitating its endocytosis into the cells.




glycolipids

Sphingolipid type	R group
Ceramide	H
Sphingomyelin	phosphocholine
Cerebroside	monosaccharide (galactose or glucose)
Globoside	two or more sugars (galactose, glucose, N-acetylglucosamine)
Ganglioside	three or more sugars including at least one sialic acid




- CCCCCCCCCCCCC=CC(O)C(O)CNC(=O)C1OC(CO)C(OS(=O)(=O)O)C1O



Sphingolipids and blood groups




- Sphingolipids serve in intercellular communication and as the antigenic determinants of the ABO blood groups.
- Some are used as receptors by viruses and bacterial toxins.




Lipoproteins



Function: transport of different types of lipids (cholesterol, cholesterol esters, phospholipids & triacylglycerols) in blood plasma.




As lipid content increases, the density decreases




Steroids



The precursor




The nucleus



Steroid nucleus





The most common steroid



Products of cholesterol




- Hormones
 - Sex hormones (androgens,. estrogens, progestins)
- Vitamin D
- Bile acids (intestinal absorption of fat)





Cholesterol esters



- A cholesterol with a fatty acid attached at (-OH) of C3



Name the molecules?




Atherosclerosis



Normal Coronary Artery with Normal blood flow




Cholestrol Deposition in Coronary Artery with Impaired blood flow



Cell membranes



- The membrane is hypothesized in a model known as the fluid mosaic model.
- Components: 45% lipid, 45% protein and 10% carbohydrate
- They exist side by side without forming some other substance of intermediate nature.




Phospholipids



- The outer: phosphatidylcholine, sphingomyelin, and glycolipids (cell recognition)
- The inner: phosphatidylethanolamine, phosphatidylserine, and phosphatidylinositol (signaling)

Cholesterol is distributed in both leaflets


Animal cells vs. plant cells vs. prokaryotic cells



Fatty acids and membrane fluidity



**Saturated
fatty acids**




**Mixture of saturated and
unsaturated fatty acids**


Cholesterol and membrane fluidity



- The presence of cholesterol and the cis unsaturated fatty acids in the membrane prevent the hydrophobic chains from packing too closely together, allowing free membrane proteins and lipid molecules to move laterally in the plane of the leaflet making the membrane a dynamic environment.
- Cholesterol can also stabilize very fluid membranes by increasing interactions between the fatty acids of phospholipids through hydrophobic interactions with the cholesterol ring structure.



Membrane proteins



Types of membrane proteins




- Peripheral proteins:
 - are associated with the exterior of membranes via noncovalent interactions
- Integral membrane proteins:
 - anchored into membrane via hydrophobic regions
- Lipid-anchored:
 - associated via a lipid group

Peripheral membrane proteins





- They are associated with membranes but do not penetrate the hydrophobic core of the membrane.
 - They can be associated with integral membrane proteins.
- They are not strongly bound to the membrane and can be removed without disrupting the membrane structure.
 - Treatment with mild detergent



Integral membrane proteins




The integral membrane proteins can be associated with the lipid bilayer in several ways.



The membrane integral domains are:
1. Single or multiple
2. α -helix (human) or β -sheet (bacteria)

Some can form channels.



Structure-function of Membranes



- Transport:
 - Membranes are impermeable barrier
 - Proteins can be carriers or channels
- Signaling
 - Protein receptors and small molecules (some can be lipids themselves)
- Catalysis
 - Enzymes