



8- Rhinoviruses, Coronaviruses, Influenza and parainfluenza viruses


Mohammad Altamimi, MD, PhD
Faculty of Medicine, Jordan University, 2024

Objectives

- Discuss the structure, properties, epidemiology, clinical presentation, laboratory diagnosis and treatment of the following viruses:
 1. Rhinoviruses
 2. Corona viruses
 3. Influenza Virus
 4. Parainfluenza viruses
 5. Respiratory syncytial virus (RSV)

Anatomy of Respiratory Tract


- The upper respiratory tract:
 - Nasal cavity, sinuses, pharynx, and larynx
 - Infections are fairly common.
 - Usually nothing more than an irritation
- The lower respiratory tract:
 - Lungs and bronchi
 - Infections are more dangerous.
 - Can be very difficult to treat
- The most accessible system in the body:
Breathing brings in clouds of potentially infectious pathogens.



Common Viral Causes

1. Rhinoviruses
2. Corona viruses
3. Influenza Virus
4. Parainfluenza viruses
5. Respiratory syncytial virus (RSV)

1. Rhinoviruses




- Family: Picornaviridae
- Genus : Rhinovirus
- Unenveloped
- Small icosahedral particle, 20 - 30 nm in diameter
- The viral genome is ss-RNA
- More than 100 types
- They replicate in the nasopharynx
- Shed in large amounts in nasal secretion
- The optimal temperature for their replication is 33-35 c
- Do not efficiently replicate at body temperature

Rhinitis (common cold)





- Rhinoviruses are responsible for 30-50% of common colds, coronaviruses 10-30%.
- Common cold: inflammation of the nose and throat (nasopharynx)
- Symptoms
 - Watery nasal discharge.
 - Sneezing.
 - Mild sore throat.
 - Fever is not common
- It is a highly contagious disease
- Transmitted by inhalation of respiratory droplets during sneezing and coughing and through contaminated hands
- No lab tests are usually required and no treatment is required only supportive treatment




2. Coronaviruses

- Family: Coronaviridae
- Irregular in shape
- Enveloped with club-shaped glycoprotein spikes
- Spike protein is important for attachment and immunity
- Helical nucleocapsid
- The viral genome is ss-RNA




Corona Diseases

- Coronaviruses can cause common cold, SARS-CoV-1 and MERS-CoV, and recently COVID-19
- COVID-19 disease caused by SARS-CoV-2
- Transmission by air droplets and direct contact
- Symptoms: Fever or chills, cold-like symptoms dry cough, shortness of breath, loss of taste or smell, fatigue, headache, vomiting, diarrhea
- Complications: acute respiratory distress syndrome
- Diagnosis: RT-PCR, chest x ray and CT-scan
- Treatment: debatable and none specific
- Vaccinations are available



6. Influenza Virus

- Influenza virus is an orthomyxovirus
- Segmented ssRNA virus
- Spherical/ovoid, 80-120 nm diameter
- The outer surface of the particle consists of a lipid envelope from which project prominent glycoprotein spikes of two types
 1. **Haemagglutinin (HA):** Can agglutinate RBCs used for viral attachment and fusion, and it elicits neutralizing protective antibody responses
 2. **Neuraminidase (NA):** Enzyme that uses neuraminic (sialic) acid as a substrate. Important in releasing mature virus from cells



Types

- Three types: A, B, and C
 - Type A undergoes antigenic shift and drift. This group is the cause of epidemics and pandemics and has an avian intermediate host (IH)
 - Type B undergoes antigenic drift only. This group causes epidemics and has no IH
 - Type C is relatively stable. This group does not cause epidemics and causes mild disease
- Subtypes:
 - According to antigenicity of HA and NA, influenza virus is divided into subtypes such as HnNm(H₁N₂, et al)

Antigenic shift and drift

- Antigenic shift:
 - Reassortment of genes is a common feature of Influenza A, but not B or C
 - When two different "A" viruses infect the same cell, their RNA segments can become mixed during replication
 - New viruses produced in this way may survive due to a selective advantage within the population
 - 1918 H₁N₁: "Spanish Influenza" 20-40 million deaths
 - 1957 H₂N₂: "Asian Flu" 1-2 million deaths
 - 1968 H₃N₂: "Hong Kong Flu" 700,000 deaths
- Antigenic drift:
 - Constant mutations in the RNA of influenza which lead to polypeptide mutations
 - Changes are less dramatic than those induced by Shift


Influenza (Flu)



- Can occur as pandemics due to antigenic shifts or Epidemics through antigenic drifts or sporadic cases
- Transmission is by respiratory droplets
- Occur more frequently in the winter
- Symptoms: Fever, headache, myalgia, cough and rhinitis
- Diagnosis: nasopharyngeal aspirates, throat, and nasal swabs are normally used for antigen detection, RT-PCR for viral RNA, Virus Isolation, and Serology
- Neuraminidase inhibitors (Tamiflu) - are now the drugs of choice.
- Vaccines are available annually with efficacy of 40-50%


3. Parainfluenza viruses

- Family: Paramyxoviridae
- Large, 150-300 nm in diameter
- Pleomorphic with helical nucleocapsid
- Enveloped with two glycoprotein spikes, HN and F
 - The HN has both hemagglutinine and neuraminidase activities used for attachment
 - The F (fusion), mediates cell entry by the fusion process
- The viral genome is ss-RNA
- 5 subtypes: 1, 2, 3, 4a and 4b
- Transmission: respiratory droplets, winter months.
- It occurs in children (below 3 years).




Croup

- Acute inflammation of the larynx and trachea in infants and young children characterized by swelling of the epithelial cells lining the airway, so that the airway narrows, and breathing becomes difficult
- Symptoms: afebrile, early runny nose, harsh cough, inspiratory stridor, and hoarse voice. Symptoms subside within 1 or 2 days.
- Diagnosis: clinical diagnosis is the rule. Lab: Detection of Antigen, Virus Isolation, and Serology
- Treatment:
 - No specific antiviral chemotherapy is available.
 - Severe cases should be admitted and placed in oxygen tents.
 - Severe respiratory obstruction may require endotracheal intubation and tracheotomy
 - No vaccine is available.




4. Respiratory syncytial virus (RSV)

- Family: Paramyxoviridae
- Large, 150-300 nm
- Pleomorphic, helical nucleocapsid
- Enveloped with two glycoprotein spikes:
 - The G protein, lacks hemagglutinins and neuraminidase activities. Attachment protein
 - The F, Mediates cell entry, by the fusion process
- The viral genome is ss-RNA
- Most common cause of severe lower respiratory tract disease in infants, responsible for 50-90% of Bronchiolitis and 5-40% of Bronchopneumonia. In older children and adults, the symptoms are much milder.



Bronchiolitis



- Inflammation of the bronchioles in infants and young children. Bronchioles become inflamed, edematous and obstructed by mucous .
- Respiratory syncytial virus (RSV) and parainfluenza virus type 3 are the major cause of bronchiolitis in infants.
- Symptoms:
 - Usually preceded by URT symptoms.
 - Expiratory obstruction.
 - Expiratory wheezing.
 - Respiratory distress (difficult & labored breathing).
 - Hypoxia and cyanosis
- Most cases are mild, recover completely & do not require hospitalization. Increasing respiratory distress, cyanosis, fatigue or dehydration are indication for hospitalization.
- Diagnosis is usually clinical and no specific treatment and no vaccination is available.

Viral pneumonia

- Inflammation of the lung and alveoli
- RSV and parainfluenza virus type-3 are the major cause of infantile pneumonia
- Symptoms: usually preceded by the URT symptoms
 - Fever
 - Chills
 - Pharyngitis
 - Cough
 - Shortness of breath
 - Muscle aches
 - Fatigue
 - Chest pain
- Prognosis: Most cases are mild and get better without treatment
- Some cases are more serious and require hospitalization
- Complications: Respiratory failure and heart failure

