

Past Papers

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ



MID – Lecture 1 to 3

# Metabolism

﴿ وَإِن تَتَوَلَّوْا يَسْتَبَدِلْ قَوْمًا غَيْرَكُمْ ثُمَّ لَا يَكُونُوا أَمْثَلَكُمْ ﴾

اللهم استعملنا ولا تستبدلنا

Written by:

- Ibraheem Samarah
- Ahmad Rami



First we will start with ( ) past Qs

then there will be ( ) test bank Qs

( all Qs will be by default past unless it  
is written to be test bank )

Q1 : If a reaction has negative  $\Delta G$  then it has to be:

A-Exergonic

B-Exothermic

C-Endothermic

D-Endergonic

E-non of the above

Answer : A

Q2 :Measure the change in the disorder of reactants and products is?

A-Delta G

B-Delta H

C-Delta S

D-Delta T

E-Delta G°

Answer : C

Q3 :Which of the following that predict whether reactions is spontaneous or not:

A-Delta  $G^\circ$

B-Delta G

C-Delta H

D-Delta E

E-Delta  $E^\circ$

Answer : B

Q 4:Delta G represents energy changes at constant temperature, pressure and proton concentration:

A-True

B-False

C-Can't be known

D-Depends on the conditions

Answer : B

Q 5:  $\Delta G = \Delta G^\circ$  , when:

A-  $R=0$

B-  $[\text{reactant}]=0$

C-  $[B]/[A]=0$

D-  $\ln [B]/[A]=1$

E-  $[B]/[A]=1$

Answer : E

Q 6: Which of the following pair is NOT true:

A- positive  $\Delta G \rightarrow$  endergonic

B- negative  $\Delta G \rightarrow$  exergonic

C-  $\Delta G = \text{Zero} \rightarrow$  equilibrium and concentration ( $[A] = [B]$ ) are equal

D-  $\Delta G$  = don't measure fast of reaction


E- exergonic reaction is favorable

Answer : C



Q7 :3-Assuming Q is the ratio of product to reactant concentrations; which of the following graph points represents accurately the equilibrium point?

- A. Point A
- B. Point C
- C. Point D
- D. Point B
- E. Cannot be known from the information given, so can be any of the above points



Answer : b

Q8 :Which one of the following reaction would you expect to be exergonic?

- a. Decarboxylation
- b. Condensation
- c. Transamination
- d. Carboxylation
- e. Phosphorylation

Answer : a

Q9 :A reaction with(-632 ) delta g, is it endergonic or exergonic reaction? And how will the addition of an enzyme affect delta g?

- A. Endergonic reaction, will not affect delta G
- B. Exergonic, will reduce activation energy
- C. exergonic reaction, will not affect delta g
- D. Endergonic, will increase activation energy
- E. Exergonic, will increase activation energy

Answer : C

Q 10: reaction has  $\Delta G^\circ > 0$ , what do you expect the value of  $K_{eq}$  ?

- A.  $K_{eq} > 1$
- B.  $K_{eq} = 0$
- C.  $K_{eq} < 1$
- D.  $K_{eq} > 0$
- E.  $K_{eq} = 1$

Answer :C

Q11 :What is the standard free energy of the reaction if  $\Delta E^\circ = -10$  mvolt, 2 electron transported, Faraday constant=23 Kcal/volt?

- A) 0.46 kcal
- B) -0.46 kcal
- C) 4.6 kcal
- D) -4.6 kcal
- E) 0.046 kcal

Answer : A

Q12 :ATP is the energy molecule of the cell because:

- A) it is the only energy molecule in the body
- B) it has 3 phosphate groups
- C) it has an intermediate energy value
- D) it isn't present in all cells
- E) C+D

Answer : C

Q 13: In experiment electrons transferred = 4 and  $\Delta E^\circ = 10$  mV calculate  $\Delta G^\circ = ?$

- a) 0.95
- b) -0.92
- c) 0.92
- d) -.092
- e) 0.092

Answer : b

Q 14:if you have the following rxns and their delta G values at standard conditions



ATP  $\rightarrow$  ADP + P<sub>i</sub>.....  $\Delta G^0 = -30.5$  The value of  $\Delta G$  at standard conditions for the following RXN equals:  $A + B + ADP \rightarrow C + ATP$

a) -73.5

b) +73.5

c) -12.5

d) +12.5

e) we can't find it out unless we have Keq

Answer : c



Q15 :If enthalpy change( $\Delta H^\circ$ ) for a reaction is zero, then  $\Delta G^\circ$  equals to :

- a)  $-T\Delta S^\circ$
- b)  $T\Delta S^\circ$
- c)  $-\Delta H^\circ$
- d)  $\ln k_{eq}$
- e)  $-\ln k_{eq}$

Answer : a

Q 16:  $\Delta G^\circ$  is defined as the :

- a) Residual energy present in the reactants at equilibrium
- b) Residual energy present in the products at equilibrium
- c) Difference in the residual energy of reactants and products at equilibrium
- d) Energy required or released to reach equilibrium when  $[\text{products}] = [\text{reactants}] = 1$
- e) Residual energy present in the products and reactants

Answer : d

Q 17: For a reaction if  $\Delta G^\circ$  is positive, then:

- a) The products will be favored
- b) The reactants will be favored
- c) The concentration of the reactants and products will be equal
- d) All of the reactant will be converted to products
- e) a+d

Answer : b

Q 18: If  $\Delta G^\circ$  of the reaction  $A \rightarrow B$  is  $-40\text{kJ/mol}$  under standard conditions then the reaction:

- a) Will never reach equilibrium
- b) Will not occur spontaneously
- c) Will proceed at a rapid rate
- d) Will proceed from left to right spontaneously
- e)  $K_{eq} < 1$

Answer : d

Q 19: Which of the following statements is true ?

- a) The reaction tends to go in the forward direction if  $\Delta G$  is large and positive
- b) The reaction tends to move in the backward direction if  $\Delta G$  is large and negative
- c) The system is at equilibrium if  $\Delta G = 0$
- d) The reaction tends to move in the backward direction if  $\Delta G_0$  is large and positive
- e) The reaction tends to go in the forward direction if  $\Delta G_0$  is large and negative

Answer : C

Q20 :The standard free energy change for a reaction in which A and B are converted to C and D is 0.4. The reaction was started by mixing 1 mmoles of each reactants and products. When the reaction reaches equilibrium, you expect that the molar concentration of:

- a. A is greater than B.
- b. A is larger than D.
- c. A less or equal to C.
- d. A and C are equal.
- e. A is larger or equal to D.

Answer : b

Q21 :The hydrolysis reaction of Glucose 6-phosphate is produces 3.3 kcal per mole under standard conditions. Calculate the standard free energy reaction for the synthesis of glucose 6-phosphate from ATP and glucose:

- a. -4.0
- b. -3.3
- c. -10.6
- d. 10.6
- e. +3.3

Answer : A

Q22 :if a non-spontaneous reaction accompanied by an increase in enthalpy , what do expect delta g:

- a. this reaction must be endothermic
- b. heat is liberated from reaction
- c. the rate of reaction is high
- d. I can't determine delta G
- e. this reaction must be exothermic

Answer : A



Q23 :A reaction has a positive delta G note, one statement is correct:

- a. This reaction will not happen in a cell.
- b. It could happen if coupled with an endergonic reaction.
- c. It can happen when changing the concentration of the reactants and the product.
- d.  $\ln K_{eq} < 1$
- e. c+d

Answer : D

Q24 :If you know that, delta E for these reactions are  $\text{NAD}^+/\text{NADH} = -0.32$ ,  $\text{pyruvate}/\text{lactate} E = -0.19$ , choose the correct statement:

- a) pyruvate/lactate is the stronger oxidizing agent.
- b)  $\text{NAD}^+/\text{NADH}$  is the stronger oxidizing agent.
- c) Pyruvate/lactate is the stronger reducing agent.
- d)  $\text{NAD}^+$  is higher tendency to gain electrons than pyruvate
- e) C+B

Answer : a

Q25 :Which one of the following cannot be a mechanism used in the body to overcome an endergonic reaction?

- a. Reaction coupling
- b. Increased substrate concentration
- c. Low intermediate concentration
- d. Decreased product concentration
- e. none of the above

Answer : C

Q 26: If you knew that the conversion of oxaloacetate to malate has a  $\Delta G$  value of +32 KJ/mol, which of the following is true:

- a. it will move slower.
- b. it will not happen in the cell.
- c. It may occur in the cell with specific concentrations for the reactant and products.
- d. It could happen if coupled with an endergonic reaction.
- e. None of the above.

Answer : C

Q 27: Which of the following concentrations of ATP and ADP are the most suitable for the human body?

	ATP	ADP	Pi
A	2	14.2	10
B	5	10	25
C	5	0.2	10

- a) A
- b) B
- c) C

Answer : C

Q28 :All of the following regarding thermodynamics are INCORRECT, except:

- a. If  $\Delta G < 0$ , reaction is spontaneous and releases energy
- b. If  $\Delta G < 0$ , reaction is spontaneous and consumes energy
- c. If  $\Delta G > 0$ , reaction is spontaneous and consumes energy
- d. If  $\Delta G > 0$ , reaction is non-spontaneous and releases energy
- e. a+b

Answer : A

Q29 :The equilibrium constant ( $K_{eq}$ ) depends on which of the following?

- a) Concentration of reactants
- b) Concentration of products
- c) Pressure
- d) Temperature
- e) Gibbs free energy

Answer : d

# Test bank Qs



**Q30** : Which of the following statements regarding equilibrium is false?

- A. Reactions with an equilibrium constant  $> 1$  favor products
- B. Concentrations of products and reactants is not necessarily equal
- C. Ratio of products to reactants is constantly increasing
- D. Rate of forward reaction is equal to rate of backward reaction
- E. all of the above

Answer : C

**Q31** : What happens to  $\Delta G$  when reactant concentrations increase significantly at equilibrium?

- A. It becomes more positive
- B. It becomes more negative
- C. Stays the same value
- D. Nears zero
- E. Not enough information provided

Answer : B

Q32 : In what cellular process is CTP primarily involved ?

- A. Carbohydrate Synthesis
- B. Combining Sugars
- C. Protein Synthesis
- D. Lipid Synthesis
- E. Phosphorylation

Answer : D

Q33 : What happens to the free energy released from ATP hydrolysis in an energy-coupled reaction?

- A. Stored in the products of the reaction
- B. Converted into entropy
- C. Used to drive a spontaneous reaction
- D. Drives an endergonic reaction
- E. Used to increase cellular temperature

Answer : D

Q34 : Which molecule is most often used alongside ATP to couple energy in protein synthesis?

A.CTP

B.UTP

C.GTP

D.FADH<sub>2</sub>

E.UMP

Answer : C

Q35 :How does  $\Delta E^\circ$  relate to the spontaneity of a redox reaction?

- A. A positive  $\Delta E^\circ$  means the reaction is spontaneous
- B. A negative  $\Delta E^\circ$  means the reaction is spontaneous
- C.  $\Delta E^\circ=0$  means reaction is at equilibrium
- D.  $\Delta E^\circ$  is not related to the spontaneity of the reaction
- E. All are correct

Answer : A

Q36 : Calculate the  $\Delta G^\circ$  for a reaction where 3 moles of electrons are transferred and the  $\Delta E^\circ$  is  $-0.2$  V (Faraday constant  $F=23.06$  kcal/volt).

- A. 13.84 kcal/mol
- B. 27.67 kcal/mol
- C. -13.84 kcal/mole
- D. -27.67 kcal/mol
- E. Can't be determined from information

Answer : A

Q37 : What is a coenzyme A molecule primarily used for?

- A. ATP synthesis
- B. Acetyl group transfer
- C. Protein Synthesis
- D. Carbohydrate storage
- E. Lipid Synthesis

Answer : B



Q38 : What is the Gibbs free energy change  $\Delta G$  for ATP hydrolysis in standard conditions?

- A. -7.5 kcal/mol
- B. -3.4 kcal/mol
- C. -7.3 kcal/mol
- D. +3.4 kcal/mol
- E. 14.6 kcal/mol

Answer : C

Q39 : What does the term “thermogenesis” refer to?

- A. Energy production for ATP synthesis
- B. Energy expended for heat generation
- C. Energy required for muscle contraction
- D. Energy lost in metabolic pathways
- E. Energy stored in fats

Answer : B

Q40 : When does a reaction reach equilibrium?

- A. When reactants are completely consumed
- B. When the rate of forward and reverse reactions are equal
- C. When the concentration of reactants equals that of products
- D. When the reaction is the irreversible
- E.  $\Delta G < 0$

Answer : B

Q41: Which of the following statements describes the relationship between  $\Delta G$  and  $\Delta G^\circ$  in cellular conditions?

- A)  $\Delta G$  equals  $\Delta G^\circ$  under all conditions
- B)  $\Delta G$  represents the free energy change under standard conditions, while  $\Delta G^\circ$  accounts for physiological conditions
- C)  $\Delta G$  is constant regardless of product and reactant concentrations
- D)  $\Delta G$  is related to  $\Delta G^\circ$  through the equation:  $\Delta G - \Delta G^\circ = RT \ln K_{eq}$
- E)  $\Delta G$  is always positive for anabolic reactions

Answer : D

Q42: A reaction has a  $\Delta G^\circ$  of +3.4 kcal/mol. Which of the following changes could allow this reaction to proceed spontaneously in a cell?

- A. Decreasing the concentration of reactants
- B. Increasing the concentration of products
- C. Coupling the reaction to ATP hydrolysis
- D. Lowering the reaction temperature
- E. Increasing the reaction activation energy

Answer : C

Q43: All of the following are characteristics of exergonic reactions except:

- A. They have a negative  $\Delta G$
- B. They release free energy to the system
- C. They are spontaneous under standard conditions
- D. They require an input of energy to proceed
- E. They are often coupled with endergonic reactions in metabolism

Answer : D

Q44: All of the following are true about thermogenesis except:

- A. Thermogenesis refers to the production of heat as a by-product of metabolic reactions.
- B. Shivering thermogenesis involves muscle contractions that generate heat
- C. Non-shivering thermogenesis is mainly associated with brown adipose tissue
- D. Non-shivering thermogenesis is an ATP-consuming process that occurs in response to cold temperatures.
- E. Thermogenesis helps maintain body temperature, particularly in cold environments

Answer : D

Q45: A reaction has a  $\Delta G^\circ$  of  $-2.5$  kcal/mol, and the ratio of reactants to products 10:1. What happens to the  $\Delta G$  value of the reaction?

- A.  $\Delta G$  becomes more negative, reaction favors forward direction
- B.  $\Delta G$  becomes more positive, reaction becomes less spontaneous
- C.  $\Delta G$  stays the same as  $\Delta G^\circ$
- D.  $\Delta G$  becomes zero, thus reaction reaches equilibrium
- E.  $\Delta G$  becomes more positive, reaction favors backward direction

Answer : A



Q46: Which of the following statements is true about the relationship between  $\Delta G$ ,  $\Delta G^\circ$ , and  $K_{eq}$ ?

- A.  $\Delta G = \Delta G^\circ$  when  $K_{eq}=0$
- B. When  $\Delta G$  is negative, reaction is always spontaneous under standard conditions
- C. When  $\Delta G^\circ$  is positive, reaction is always non-spontaneous under standard conditions
- D.  $\Delta G$  depends only on the concentration of reactants
- E.  $\Delta G = \Delta G^\circ$  only when the temperature is  $25^\circ\text{C}$

Answer : C

Q47: All of the following statements is true about  $\Delta G$  except:

- A.  $\Delta G$  can determine whether reaction is spontaneous or not
- B.  $\Delta G$  is not affected by the reaction mechanism or pathway
- C.  $\Delta G$  can't be altered by enzymes catalyzing the reaction
- D.  $\Delta G$  only depends on the initial and final states of the reaction
- E. A reaction with  $\Delta G=1$  is at equilibrium

Answer : E

For any feedback, scan the code or click on it.



Corrections from previous versions:

Versions	Question #	Before Correction	After Correction
V1 → V2	21	E	A
	29	E	D
	31	A	B
	44	A	D
	45	E	A
V2 → V3			