

بسم الله الرحمن الرحيم



FINAL | Lecture #2


Axilla & Brachial Plexus

Written by: Moneeb Arabiat
Mazen Alnashash

Reviewed by: Laith Joudeh

وَأِنْ تَتَوَلَّوْا يَسْتَبَدِلْ قَوْمًا غَيْرَكُمْ ثُمَّ لَا يَكُونُوا أَمْثَلَكُمْ

اللهم استعملنا ولا تستبدلنا



ANATOMY




Color Code: Slides + Dr. doesn't mention Slides + Dr. mentions Extra from Dr.

The Axilla


﴿ وَتَوَكَّلْ عَلَى الْحَيِّ الَّذِي لَا يَمُوتُ وَسَبِّحْ بِحَمْدِهِ ۚ وَكَفَىٰ بِهِ بِذُنُوبِ عِبَادِهِ خَبِيرًا ﴾
[الفرقان: 58]

The Axilla

- The axilla, or armpit, is a pyramid-shaped space between the upper part of the arm and the side of the chest
- It forms an important passage for nerves, blood, and lymph vessels as they travel from the root of the neck to the upper limb.
- The upper end of the axilla, or apex, is directed into the root of the neck
- is bounded in front by the clavicle, behind by the upper border of the scapula, and medially by the outer border of the first rib




- The lower end, or base, is bounded in front by the anterior axillary fold (formed by the lower border of the pectoralis major muscle),
- behind by the posterior axillary fold (formed by the tendon of latissimus dorsi and the teres major muscle),
- and medially by the chest wall



Walls of the Axilla

- Anterior wall: By the pectoralis major, subclavius, and pectoralis minor muscles **along with the clavipectoral fascia, which surrounds the pectoralis minor and extends to the floor of the axilla as the suspensory ligament.**
- Posterior wall: By the subscapularis, latissimus dorsi, and teres major muscles from above down
- Medial wall: By the upper four or five ribs and the intercostal spaces covered by the serratus anterior muscle




anterior wall




posterior wall

- Lateral wall: By the coracobrachialis and biceps muscles in the bicipital groove of the humerus
- The base is formed by the skin stretching between the anterior and posterior walls
- The axilla contains the principal vessels and nerves to the upper limb and many lymph nodes
- The axilla contains the axillary artery, axillary vein, axillary lymph nodes, and the cords of the brachial plexus.



The Axilla




© Elsevier. Drake et al: Gray's Anatomy for Students - www.studentconsult.com

Axillary artery: Begins at the outer border of the first rib as a continuation of the subclavian artery and ends at the lower border of the teres major muscle, where it continues as the brachial artery.

Clavipectoral Fascia

- The clavipectoral fascia is a strong sheet of connective tissue that is attached above to the clavicle
- Below, it splits to enclose the pectoralis minor muscle
- then continues downward as the suspensory ligament of the axilla and joins the fascial floor of the armpit.
- **Structures travel between the axilla and the anterior wall of the axilla by passing through the clavipectoral fascia**
- either between the pectoralis minor and subclavius muscles or inferior to the pectoralis minor muscle.
- **Important structures that pass between the subclavius and pectoralis minor muscles include the cephalic vein, the thoraco-acromial artery, and the lateral pectoral nerve & lymphatic vessels**

- Clavi: begins in the clavicle
- Pectoral: surrounding pectoralis minor (begins in the 2nd 3rd 4th ribs and ends in the coracoid process)



Contents of the Axilla

- The axilla contains the axillary artery and its branches, which supply blood to the upper limb
- the axillary vein and its tributaries, which drain blood from the upper limb
- lymph vessels and lymph nodes, which drain lymph from the upper limb and the breast and from the skin of the trunk, down as far as the level of the umbilicus
- Lying among these structures in the axilla is an important nerve plexus, the brachial plexus, which innervates the upper limb
- These structures are embedded in fat.


Introduction to the Nervous System

سبحان الله والحمد لله ولا اله الا الله والله اكبر

Nervous System

- The nervous system is divided into two main parts: the central nervous system, which consists of the brain and spinal cord
- the peripheral nervous system, which consists of 12 pairs of cranial nerves and 31 pairs of spinal nerves and their associated ganglia.
- Functionally, the nervous system can be further divided into the somatic nervous system, which controls voluntary activities
- and the autonomic nervous system, which controls involuntary activities.

- total of 31 pairs of spinal nerves leave the spinal cord and pass through intervertebral foramina in the vertebral column
- The spinal nerves are named according to the region of the vertebral column with which they are associated:
- 8 cervical, 12 thoracic, 5 lumbar, 5 sacral, and 1 coccygeal
- Each spinal nerve is connected to the spinal cord by two roots: the anterior root and the posterior root
- The anterior root (ventral root) consists of bundles of nerve fibers carrying nerve impulses away from the central nervous system. Such nerve fibers are called efferent fibers.




- Those efferent fibers that go to skeletal muscle and cause them to contract are called motor fibers
- Their cells of origin lie in the anterior gray horn of the spinal cord.

Course of the motor nerve:


The motor nerve originates from the anterior horn → travels through the anterior (ventral) root → to the spinal nerve → and then to the muscles.

Course of the sensory nerve:


The sensory nerve originates from the skin → travels through the spinal nerve → enters the posterior (dorsal) root → passes through the dorsal root ganglion → and then to the posterior horn.



- The posterior root (dorsal root) consists of bundles of nerve fibers that carry impulses to the central nervous system and are called afferent fibers
- Because these fibers are concerned with conveying information about sensations of touch, pain, temperature, and vibrations, they are called sensory fibers
- The cell bodies of these nerve fibers are situated in a swelling on the posterior root called the posterior root ganglion




- At each intervertebral foramen, the anterior and posterior roots unite to form a spinal nerve
- Here, the motor and sensory fibers become mixed together, so that a spinal nerve is made up of a mixture of motor and sensory fibers
- On emerging from the foramen, the spinal nerve divides into a large anterior ramus and a smaller posterior ramus
- The posterior ramus passes posteriorly around the vertebral column to supply the muscles and skin of the back
- The anterior ramus continues anteriorly to supply the muscles and skin over the anterolateral body wall and all the muscles and skin of the limbs




Plexuses

- At the root of the limbs, the anterior rami join one another to form complicated nerve plexuses
- The cervical and brachial plexuses are found at the root of the upper limbs, and the lumbar and sacral plexuses are found at the root of the lower limbs.



Autonomic Nervous System

- The autonomic nervous system is the part of the nervous system concerned with the innervation of involuntary structures such as the heart, smooth muscle, and glands throughout the body
- distributed throughout the central and peripheral nervous system
- The autonomic system may be divided into two parts the sympathetic and the parasympathetic and both parts have afferent and efferent nerve fibers
- The activities of the sympathetic part of the autonomic system prepare the body for an emergency




The Brachial Plexus

وَلَا تُخْزِنِي يَوْمَ يُبْعَثُونَ (87) يَوْمَ لَا يَنْفَعُ مَالٌ وَلَا بَنُونَ (88) إِلَّا مَنْ أَتَى اللَّهَ بِقَلْبٍ سَلِيمٍ


Brachial Plexus

- The nerves entering the upper limb provide the following important functions:
 1. sensory innervation to the skin and deep structures, such as the joints
 2. motor innervation to the muscles
 3. influence over the diameters of the blood vessels by the sympathetic vasomotor nerves
 4. and sympathetic secretomotor supply to the sweat glands.

- At the root of the neck, the nerves form a complicated plexus called the brachial plexus
- The brachial plexus is formed in the posterior triangle of the neck by the union of the anterior rami of the fifth, sixth, seventh, and eighth cervical and the first thoracic spinal nerves
- The plexus can be divided into roots, trunks, divisions, and cords
- The roots (value of the nerve) of C5 and 6 unite to form the upper trunk the root of C7 continues as the middle trunk and the roots of C8 and T1 unite to form the lower trunk




- Each trunk then divides into anterior and posterior divisions
- The anterior divisions of the upper and middle trunks unite to form the lateral cord
- the anterior division of the lower trunk continues as the medial cord
- and the posterior divisions of all three trunks join to form the posterior cord
- The roots, trunks, and divisions of the brachial plexus reside in the lower part of the posterior triangle of the neck
- The cords become arranged around the axillary artery in the axilla
- the brachial plexus and the axillary artery and vein are enclosed in the axillary sheath.



Cords of the Brachial Plexus


- All three cords of the brachial plexus lie above and lateral to the first part of the axillary artery
- The medial cord crosses behind the artery to reach the medial side of the second part of the artery
- The posterior cord lies behind the second part of the artery,
- and the lateral cord lies on the lateral side of the second part of the artery
- Thus, the cords of the plexus have the relationship to the second part of the axillary artery that is indicated by their names.
- Most branches of the cords that form the main nerve trunks of the upper limb continue this relationship to the artery in its third part




The branches of the different parts of the brachial plexus

- Roots
 - Dorsal scapular nerve (C5)
 - Long thoracic nerve (C5, 6, and 7)
- Upper trunk
 - Nerve to subclavius (C5 and 6)
 - Suprascapular nerve (supplies the supraspinatus and infraspinatus muscles)
- Lateral cord:
 - Lateral pectoral nerve
 - Musculocutaneous nerve
 - Lateral root of median nerve

Middle and Lower trunk do not give immediate branches




- Medial cord:
 - Medial pectoral nerve
 - Medial cutaneous nerve of arm and medial cutaneous nerve of forearm
 - Ulnar nerve
 - Medial root of median nerve
- Posterior cord:
 - Upper and lower subscapular nerves
 - Thoracodorsal nerve
 - Axillary nerve
 - Radial nerve




Branches of the Brachial Plexus Found in the Axilla


- The nerve to the subclavius (C5 and 6) supplies the subclavius muscle
- It is important clinically because it may give a contribution (C5) to the phrenic nerve; this branch, when present, is referred to as the accessory phrenic nerve.
- The long thoracic nerve (C5, 6, and 7) arises from the roots of the brachial plexus in the neck
- enters the axilla by passing down over the lateral border of the first rib behind the axillary vessels and brachial plexus
- It descends over the lateral surface of the serratus anterior muscle, which it supplies.



- The lateral pectoral nerve arises from the lateral cord of the brachial plexus and supplies the pectoralis major muscle
- The musculocutaneous nerve arises from the lateral cord of the brachial plexus
- supplies the coracobrachialis muscle, and leaves the axilla by piercing that muscle




- The lateral root of the median nerve is the direct continuation of the lateral cord of the brachial plexus
- It is joined by the medial root to form the median nerve trunk, and this passes downward on the lateral side of the axillary artery
- The median nerve gives off no branches in the axilla
- The medial pectoral nerve arises from the medial cord of the brachial plexus
- supplies and pierces the pectoralis minor muscle, and supplies the pectoralis major muscle




- The medial cutaneous nerve of the arm (T1) arises from the medial cord of the brachial plexus
- And is joined by the intercostobrachial nerve (lateral cutaneous branch of the second intercostal nerve).
- It supplies the skin on the medial side of the arm.


- The medial cutaneous nerve of the forearm arises from the medial cord of the brachial plexus and descends in front of the axillary artery
- The ulnar nerve (C8 and T1) arises from the medial cord of the brachial plexus and descends in the interval between the axillary artery and vein
- The ulnar nerve gives off no branches in the axilla
- The medial root of the median nerve arises from the medial cord of the brachial plexus and crosses in front of the third part of the axillary artery to join the lateral root of the median nerve



- The upper and lower subscapular nerves arise from the posterior cord of the brachial plexus and supply the upper and lower parts of the subscapularis muscle
- In addition, the lower subscapular nerve supplies the teres muscle
- The thoracodorsal nerve arises from the posterior cord of the brachial plexus and runs downward to supply the latissimus dorsi muscle (Also called nerve to latissimus dorsi muscle).



- The axillary nerve is one of the terminal branches of the posterior cord of the brachial plexus
- Having given off a branch to the shoulder joint, it divides into anterior and posterior branches
- The radial nerve is the largest branch of the brachial plexus and lies behind the axillary artery
- It gives off branches to the long and medial heads of the triceps muscle and the posterior cutaneous nerve of the arm
- The latter branch is distributed to the skin on the middle of the back of the arm
- **Radial nerve supplies all extensor muscles of the hand and forearm**




Blood Vessels in The Axilla

فَلَمَّا تَرَآءَى الْجَمْعَانِ قَالَ أَصْحَابُ مُوسَى إِنَّا لَمُدْرِكُونَ (61) قَالَ كَلَّا إِنَّ مَعِيَ رَبِّي سَيَهْدِينِ (62)


Axillary Artery

- The axillary artery begins at the lateral border of the first rib as a continuation of the subclavian
- and ends at the lower border of the teres major muscle, where it continues as the brachial artery
- Throughout its course, the artery is closely related to the cords of the brachial plexus and their branches and is enclosed with them in a connective tissue sheath called the axillary sheath
- If this sheath is traced upward into the root of the neck, it is seen to be continuous with the prevertebral fascia.
- The pectoralis minor muscle crosses in front of the axillary artery and divides it into three parts




First Part of the Axillary Artery

- This extends from the lateral border of the first rib to the upper border of the pectoralis minor
- **The first part of the axillary artery gives one branch: the highest thoracic artery.**
- Relations
 - Anteriorly: The pectoralis major and the skin. The cephalic vein crosses the artery
 - Posteriorly: The long thoracic nerve (nerve to the serratus anterior)
 - Laterally: The three cords of the brachial plexus
 - Medially: The axillary vein (**MOST MEDIAL**)




Second Part of the Axillary Artery

- This lies behind the pectoralis minor muscle
- Relations
 - Anteriorly: The pectoralis minor, the pectoralis major, and the skin
 - Posteriorly: The posterior cord of the brachial plexus, the subscapularis muscle, and the shoulder joint
 - Laterally: The lateral cord of the brachial plexus
 - Medially: The medial cord of the brachial plexus and the axillary vein




Third Part of the Axillary Artery

- This extends from the lower border of the pectoralis minor to the lower border of the teres major
- Relations:
- Anteriorly: The pectoralis major for a short distance; lower down the artery it is crossed by the medial root of the median nerve
- Posteriorly: The subscapularis, the latissimus dorsi, and the teres major. The axillary and radial nerves also lie behind the artery
- Laterally: The coracobrachialis, the biceps, and the humerus. The lateral root of the median and the musculocutaneous nerves also lie on the lateral side
- Medially: The ulnar nerve, the axillary vein, and the medial cutaneous nerve of the arm




Branches of the Axillary Artery

1. 1) From the first part:
 - The highest thoracic artery is small and runs along the upper border of the pectoralis minor
2. From the second part:
 - A. The thoracoacromial artery (**trunk**) immediately divides into terminal branches.
 - B. The lateral thoracic artery runs along the lower border of the pectoralis minor




3. From the third part:

- The subscapular artery runs along the lower border of the subscapularis muscle.
- The anterior and posterior circumflex humeral arteries wind around the front and the back of the surgical neck of the humerus, respectively



Axillary Vein (Most medial in axilla)


- The axillary vein is formed at the lower border of the teres major muscle by the union of the venae comitantes of the brachial artery and the basilic vein
- It runs upward on the medial side of the axillary artery and ends at the lateral border of the first rib by becoming the subclavian vein.
- The vein receives tributaries, which correspond to the branches of the axillary artery, and the cephalic vein



Axillary Lymph Nodes

وَاجْعَلْنِي مِنْ وَرَثَةِ جَنَّةِ النَّعِيمِ

- The lymph nodes are arranged in six groups
- **All six groups of axillary lymph nodes are commonly removed during breast cancer surgery**
- 1)Anterior (pectoral) group:
 - Lying along the lower border of the pectoralis minor behind the pectoralis major
 - receive lymph vessels from the lateral quadrants of the breast and superficial vessels from the anterolateral abdominal wall




2. Posterior (subscapular) group:

- Lying in front of the subscapularis muscle
- receive superficial lymph vessels from the back, down as far as the level of the iliac crests.

3. Lateral group :

- Lying along the medial side of the axillary vein
- receive most of the lymph vessels of the upper limb (except those superficial vessels draining the lateral side)




4. Central group:

- Lying in the center of the axilla in the axillary fat
- receive lymph from the above three groups (**Anterior+posterior+lateral**)


5. Infraclavicular (deltopectoral) group:

- they are located outside the Axilla
- They lie in the groove between the deltoid and pectoralis major muscles
- receive superficial lymph vessels from the lateral side of the hand, forearm, and arm



6. Apical group :

- Lying at the apex of the axilla at the lateral border of the first rib
- receive the efferent lymph vessels from all the other axillary lymph nodes.
- The apical nodes drain into the subclavian lymph trunk
- On the left side, this trunk drains into the thoracic duct; on the right side, it drains into the right lymph trunk
- the lymph trunks may drain directly into one of the large veins at the root of the neck.



Drainage of Apical Lymph Nodes into the Subclavian Lymph Trunk

- **Right side:** Lymph drains into the **right lymphatic duct**, then to the **right venous angle**, and finally into the **right brachiocephalic vein**.
- **Left side:** Lymph drains into the **thoracic duct**, then to the **left venous angle**, and finally into the **left brachiocephalic vein**.

For any feedback, scan the code or click on it.



Corrections from previous versions:

Versions	Slide # and Place of Error	Before Correction	After Correction
V0 → V1			
V1 → V2			

رسالة من الفريق العلمي:

رَبِّ اغْفِرْ لِي وَلِوَالِدَيَّ وَلِمَن دَخَلَ بَيْتِي مُؤْمِنًا وَلِلْمُؤْمِنِينَ
وَالْمُؤْمِنَاتِ وَلَا تَزِدِ الظَّالِمِينَ إِلَّا تَبَارًا (28)

اللَّهُمَّ أَنْتَ رَبِّي، لَا إِلَهَ إِلَّا أَنْتَ
خَلَقْتَنِي وَأَنَا عَبْدُكَ، وَأَنَا عَلَى
عَهْدِكَ وَوَعْدِكَ مَا اسْتَطَعْتُ، أَعُوذُ
بِكَ مِنْ شَرِّ مَا صَنَعْتُ، أُبَوِّءُ لَكَ
بِنِعْمَتِكَ عَلَيَّ، وَأُبَوِّءُ بِذَنْبِي فَاغْفِرْ
لِي، فَإِنَّهُ لَا يَغْفِرُ الذُّنُوبَ إِلَّا أَنْتَ.