
سسم اللك الرّحون الرّحيم

Medical Immunology



Anas Abu-Humaidan M.D. Ph.D.


Lecture 2

#### Immunology introduction

Remember

- The immune system includes the role of **physical**, **cellular**, and **chemical systems** that are in place and that respond to all aspects of **foreignness**.
- The immune system targets any "foreign" object, so the first step is to **recognize** what is self and non-self.
- The second step is to **restore** homeostasis by eliminating the foreign object.
- The third step is to **remember** the invading pathogen to respond better the next time it is encountered .
- The immune system is not **only** active when danger arises, but is constantly sensing danger and is **important for normal physiology and homeostasis** similar to the cardiovascular and renal systems.


#### Immunology introduction / INNATE AND ADAPTIVE IMMUNITY



• Host defenses are grouped under **innate** immunity, which provides **immediate** protection against microbial invasion, and **adaptive** immunity, which develops **more slowly** and provides more specialized defense against infections

#### Cells of the immune system


- All immune cells *originate* from the bone marrow, but some migrate to other organs for maturation or long-term maintenance.
- The cells of the innate and adaptive immune system are normally present as circulating cells in the **blood** and **lymph**, as anatomically defined collections in **lymphoid organs**, and as scattered cells in **virtually all tissues**.



#### Cells of the immune system


- Phagocytes
- Mast Cells, Basophils, Eosinophils
- Antigen-Presenting Cells
- Lymphocytes

سحان المآث و تحدده سبحان الله العظیم می




### Cells of the immune system / Phagocytes

- Phagocytes including
  neutrophils and
  macrophages, are cells
  whose primary function is
  to identify, ingest, and
  destroy microbes.
- Phagocytes also communicate with other cells in ways that promote or regulate immune responses.



#### Cells of the immune system / Phagocytes/ Neutrophils

- Neutrophils, also called polymorphonuclear leukocytes, are the most abundant population of circulating white blood cells and mediate the earliest phases of inflammatory reactions.
  - The nucleus of a neutrophil is segmented into 3-5 connected lobules
  - Short lifespan, they circulate for about 6 hours, Production of neutrophils is stimulated by granulocyte colonystimulating factor (G-CSF). An adult human produces more than 1 × 10<sup>11</sup> neutrophils per day




#### Cells of the immune system / Phagocytes/ Neutrophils

- The cytoplasm contains granules of two types. The majority, called **specific granules**, are filled with enzymes such as lysozyme, collagenase, and elastase. The remainder are **azurophilic granules**, which are lysosomes containing enzymes and other microbicidals substances.
- Neutrophils may migrate to sites of infection


  within a few hours after the entry of


  microbes.

  The work Imp Sunction of them
- After entering tissues, neutrophils function for a few hours and then die.



#### Cells of the immune system / Phagocytes/ Neutrophils





Anothe special fonction

• Neutrophil extracellular traps (NETs) are networks of extracellular fibers, primarily composed of DNA from neutrophils, which bind pathogens.

ملعض


Functions include 8


1\_ Phagocytosis

2- Secreting granules that outsin digestive enzymes

2- NETS

# اللكم جلة وسلم على نبيتا فحد





Blood film with a striking absence of neutrophils, leaving only red blood cells and platelets

Neutropenia is an abnormally low concentration of neutrophils in the blood. Neutropenia has many causes and can be congenital and acquired (e.g. cancer treatment, autoimmune diseases).

Glover + Newtrapenia

سحان الله و محده سبحان الله العظيم بي


### **Febrile Neutropenia**

JP is a 34-year-old Caucasian male who is admitted to inpatient oncology service for induction chemotherapy for a recent diagnosis of <u>acute myeloid</u> leukemia (AML). His induction chemotherapy regimen consists of 7 + 3 induction chemotherapy with cytarabine and daunorubicin. He was placed on neutropenic precautions, started on appropriate **antimicrobial prophylaxis**, and a port was placed for **chemotherapy administration**. Ten days after the completion of his induction chemotherapy (day 17), he spiked a **fever of** 38.8°C (101.8°F) and complained of chills and nausea.

Z. 32 2 0 0 0 ml 2 Chemo therapy + Loukemia

Cells of the immune system / Phagocytes / Mononuclear Phagocytes

- The cells of the mononuclear phagocyte system originate from a common precursor in the bone marrow, circulate in the blood as monocytes, and mature and become activated in various tissues.
- Once monocytes enter tissues, mature and become macrophages. Macrophages in different tissues have been given special names to designate specific locations.



#### Cells of the immune system / Phagocytes/ Macrophages

ال حول و ل قوة إلّ الله العلي العظيم

- In addition to ingesting microbes, macrophages also ingest dead host cells as part of the cleaning up process after infection or sterile tissue injury.
- Activated macrophages secrete proteins, called cytokines, that bind to signalling receptors
  on other cells and thereby instruct those cells to respond in ways that contribute to host
  defence.
- Macrophages serve as APCs that display antigens to and activate T lymphocytes.
- Another important function of macrophages is to promote **repair of damaged tissues** by Secretive stimulating new blood vessel growth (angiogenesis) and synthesis of collagen-rich extracellular matrix (fibrosis).

#### Cells of the immune system / Phagocytes/ Macrophages




troit fra)

Macrophage-like cells are phylogenetically the poldest mediators of innate immunity Drosophila responds to infection by surrounding microbes with "hemocytes,"

which are similar to macrophages, and these cells phagocytose the microbes and wall off the infection.






Macrophages can acquire distinct functional capabilities, depending on the types of activating stimuli

## Cells of the immune system / Mast Cells, Basophils, Eosinophils

- All three cell types share the common feature of having cytoplasmic granules filled with various inflammatory and antimicrobial mediators.
- Another common feature of these cells is their involvement in immune responses that protect against helminths and immune responses that cause allergic diseases

## اللكم جلة وسلم على نبيتها فحد



#### Cells of the immune system / Mast Cells


أهم واحر


 Mast cells are bone marrow—derived cells that are present in the skin and mucosal epithelium and contain abundant cytoplasmic granules filled with cytokines histamine, and other mediators.

Mature mast cells are not found in the circulation but are constitutively present in healthy tissues, usually adjacent to small blood vessels and nerves.

Mast cells express <u>plasma membrane</u>
 receptors for <u>lgE and lgG</u> antibodies and are
 usually coated with these antibodies








#### Cells of the immune system / Basophils

- Basophils are blood granulocytes with many structural and functional similarities to mast cells.
- Like mast cells, basophils express IgG and IgE receptors, bind IgE, and can be triggered by antigen binding to the IgE.
- Basophils constitute less than 1% of blood leukocytes, normally not present in tissues and their importance is uncertain.
- When they participate in **allergic reactions** or **responses to parasites**, basophils release histamine and many other biologically active molecules that can contribute to inflammation.

#### Cells of the immune system / Eosinophils

- Eosinophils were first described in 1879 by Paul Ehrlich, who noted their unusual capacity to be stained by **acidophilic dyes**.
- They are released into the peripheral blood in a phenotypically mature state, and they are capable of being activated and recruited into tissues in response to appropriate stimuli, most notably the cytokine interleukin-5 (IL-5) and the eotaxin chemokines.
- Eosinophils spend only a brief time in the peripheral blood (they have a half-life of ~18 hrs) before they migrate to the thymus or gastrointestinal tract, where they reside under homeostatic conditions


## اللكم جلة دسلم على نبيتها فحد




Nature Reviews | Immunology

Cells of the immune system / Eosinophils





- Eosinophils are blood granulocytes that express cytoplasmic granules containing enzymes that are harmful to the cell walls of parasites but can also damage host tissues.
- Eosinophils are a common feature of the inflammatory response that occurs in asthma, as they are recruited to the lungs and airways by cytokines that are released from activated TH2 cells and by a range of chemokines, most notably those of the eotaxin family.
- Several lines of evidence suggest that deficiency of eosinophils is **not associated** with any characteristic abnormality.



Eosinophils: changing perspectives in health and disease https://www.nature.com/articles/nri3341

#### Cells of the immune system

اللهم على وسلَّم على سَيِّنا في وعلى ١٦ وعلمه ومهمه وعلى نا سَعِر ودالاه إى يدم الله ين

		TABLE 2–1 Normal Blood Cell Counts		
			Mean Number per Microliter	Normal Range
paper were ppings		White blood cells (leukocytes)	7400	4500-11,000
	1	Neutrophils	4400	1800-7700
	Ч	Eosinophils	200	0-450
	5	Basophils	40	0-200
	2	Lymphocytes	2500	1000-4800
	3	Monocytes	300	0-800

 Although most of these cells are found in the blood, their responses to microbes are usually localized to tissues.




FIGURE 2-1 Morphology of neutrophils, mast cells, basophils, and eosinophils. A, The light micrograph of a Wright-Giemsa-stained blood neutrophil shows the multilobed nucleus, because of which these cells are also called polymorphonuclear leukocytes, and the faint cytoplasmic granules. B. The light micrograph of a Wright-Giemsa-stained section of skin shows a mast cell (arrow) adjocent to a small blood vessel, identifiable by the red blood cell in the lumen. The cytoplasmic granules in the mast cell, which are stained purple, are filled with histamine and other mediators that act on adjacent blood vessels to promote increased blood flow and delivery of plasma proteins and laukocytes into the tissue. (Courtesy of Dr. George Murphy, Department of Pathology, Bigham and Women's Hospital, Baston, Massachusetts J. The light micrograph of a Wright-Giemsa-stained blood basophil shows the characteristic blue-staining cytoplasmic granules. (Courtesy of Dr. Jonethan Hecht, Department of Pathology, Bighem and Women's Hospital, Boston, Massachusetts.) D, The light micrograph of a Wright-Giemsa-stained blood eosinophil shows the characteristic segmented nucleus and red staining of the cytoplasmic granules.

# **Further reading:**

• Cellular and Molecular Immunology. 7th Edition.. Chapter 2. Cells and tissues of the immune system

سحان اللآه و تحدده سحان اللآم العقيم بي المتعاد لل منسوا إحوا من من المتعاد لله المرابطيي الما الما المواسية المعاد المعادية