
سم الله الرحمن الرحيم

Medical Immunology




Anas Abu-Humaidan M.D. Ph.D.

Lecture 3


### Cells of the immune system

- Phagocytes
- Mast Cells, Basophils, Eosinophils
- Antigen-Presenting Cells
- Lymphocytes

سحان المآه و تحده سبحان الله العظيم في




Cells of the immune system / Antigen presenting cells




- Antigen-presenting cells (APCs) are cell populations that are specialized to capture
  microbial and other antigens, display them to lymphocytes, and provide signals that
  stimulate the proliferation and differentiation of the lymphocytes.
- The major type of APQ that is involved in initiating T cell responses is the **dendritic cell**.
- Macrophages and B cells as well present antigens to T lymphocytes in different types of immune responses.
- APCs link responses of the innate immune system to responses of the adaptive immune system, and therefore they may be considered components of both systems.

#### Cells of the immune system / Antigen presenting cells/ Dendritic Cells

- Dendritic cells are the most important APCs for activating naive T cells, and they play major roles in innate responses to infections and in linking innate and adaptive immune responses.
- They have long membranous projections and phagocytic capabilities and are widely distributed in lymphoid tissues, mucosal epithelium, and organ parenchyma.
- In response to activation by microbes, conventional dendritic cells in skin, mucosa, and organ parenchyma become mobile, migrate to lymph nodes, and display microbial antigens to T lymphocytes.



Dendritic cells in skin




Artistic rendering of the surface of a human dendritic cell illustrating sheet-like processes that fold back onto the membrane surface.

Innate immunity/ Dendritic cells


اللهم مهل وسلّم ملى بنسِّما عسّ رعلى آنه و محمه وعلى من تَنْعِه دوالاه إلى بوم الدّين

main function



DC include two main cell types, the plasmacytoid DC (pDC) that are expert in type I interferon synthesis upon viral stimulation and the conventional DC (cDC) that are specialized in antigen capture, processing, and presentation for T-cell priming.

Cells of the immune system / Antigen-Presenting Cells/ Follicular Dendritic Cells




- Follicular dendritic cells (FDCs) are cells with membranous projections that are found intermingled in specialized collections of activated B cells, called **germinal** centers, in the lymphoid follicles of the lymph nodes, spleen, and mucosal lymphoid tissues.
- FDCs are not derived from precursors in the bone marrow and are of mesenchymal origin, they are **non migratory**.
- FDCs trap antigens complexed to antibodies or complement products and display these antigens on their surfaces for recognition by B lymphocytes.

Cells of the immune system / Antigen-Presenting Cells for Effector T Lymphocytes

لاحول ولا قَوَة الَّهُ عاللَّ العَظيم

- Macrophages present antigen to helper T lymphocytes at the sites of infection, which leads to helper T cell activation and production of molecules that further activate the macrophages. This process is important for the eradication of microbes that are ingested by the phagocytes but resist killing.
- **B cells present** antigens to helper T cells in lymph nodes and spleen, which is a key step in the cooperation of helper T cells with B cells in humoral immune responses to protein antigens.




# Major histocompitability complex I

- Their function is to display peptide fragments of proteins from within the cell to cytotoxic T cells; this will trigger an immediate response from the immune system against a particular non-self antigen displayed with the help of an MHC class I protein.
- The proteasome is a macromolecule that consists of 28 subunits, of which half affect proteolytic activity.

All nucleafeel b

 Cells constantly break down proteins and present them on MHC I.


سبحان الله و محده سبحان الله العظيم



# Major histocompitability complex II

لاحول ولا قَوَة الله عاللت العظيم

- Normally found only on professional antigenpresenting cells such as dendritic cells,
   Macrophages, B cells and some endothelial cells.
- The antigens presented by class II peptides are derived from extracellular proteins (not cytosolic as in MHC class I).



Page 10 of 23






FIGURE 2–11. Antigen processing and presentation. A. Antigens originating in the cytoplasm are digested by the proteasome to peptides. The peptides are bound to the MHC class I molecules in the endoplasmic reticulum (ER) and transported to the surface for presentation. B. Antigens originating outside the cell are endocytosed and digested in the phagolysosome. The digested peptides are bound to MHC class II molecules in the ER and transported to the surface for presentation. MHC, major histocompatibility complex.



**Dendritic cell** presents antigens to **lymphocytes** 

### Cells of the immune system


- Phagocytes
- Mast Cells, Basophils, Eosinophils
- Antigen-Presenting Cells
- Lymphocytes



Cells of the immune system / Lymphocytes / Natural killer (NK) cells

- Natural killer (NK) cells are lymphocytes
  distinct from T and B cells that play important
  roles in innate immune responses mainly
  against intracellular viruses and bacteria.
- The term natural killer derives from the fact that these cells are capable of performing their killing function without a need for clonal expansion and differentiation.
- Most NK cells express inhibitory receptors that recognize class I major histocompatibility complex (MHC) molecules, which are cell surface proteins normally expressed on almost all healthy cells in the body

سبحان الله و بعدد ۵ سبحان الله العظم ۶




بجعن الفيورسان عندها لهريقة للتعني والعرون من جار الهناعة عن المريقة من جلال الهناعة عن المريقة منع وكلال عن المريقة منع وكلال عن المريقة منع وكلال المريقة وكلال المريقة والمريقة وكلال المريقة وكلال المر

و رحن الحسم ييون يستجييل لفاي العِروسانَ ، ليشَ >

لاً نه عدد الله الله الله الله بعل في انظهان الطبيعية على تشابيط الاستجابة المنافسية

لبعض المخلا يا يكي متعربهن إي

## Cells of the immune system / Lymphocytes / Natural killer (NK) cells



Cells of the immune system / Lymphocytes

- Lymphocytes consist of distinct subsets that are different in their functions and protein products, but are morphologically similar.
- B (Bursa of Fabricius) lymphocytes originate in the bone marrow and early maturation occurs there.
   Also, T (Thymus) lymphocytes originate in the bone marrow, but mature in the thymus.
- Membrane proteins are used as phenotypic markers
   to distinguish distinct populations of lymphocytes





FIGURE 2-7 Morphology of lymphocytes. A, Light micrograph of a lymphocyte in a peripheral blood smear. (Courtesy of Jean Shafer, Department of Pathology, University of California, San Diego. Copyright 1995-2008, Carden Jennings Publishing Co., Ltd.) B, Electron micrograph of a small lymphocyte. (Courtesy of Dr. Noel Weidner, Department of Pathology, University of California, San Diego.) C, Light micrograph of a large lymphocyte (lymphoblast). (Courtesy of Jean Shafer, Department of Pathology, University of California, San Diego. Copyright 1995-2008, Carden Jennings Publishing Co., Ltd.) D, Electron micrograph of a large lymphocyte (lymphoblast). (From Fawcett DW. Bloom and Fawcett: A Textbook of Histology, 12th ed. Chapman & Hall, New York, 1994. With kind permission of Springer Science and Business Media.)

Cells of the immune system / Cluster of differentiation

- The cluster of differentiation (CD) is a protocol used for the identification and investigation of cell surface molecules providing targets for immunophenotyping of cells.
- In terms of physiology, CD molecules can act in numerous ways, often acting as receptors or ligands important to the cell and some function as adhesion molecules.



CD34




ك إله إن ش

Cells of the immune system / Lymphocytes/ B-lymphocytes

- B-lymphocytes are the only cells capable of producing antibodies.
- They recognize extracellular (including cell surface) antigens and differentiate into antibody-secreting plasma cells, thus functioning as the mediators of humoral immunity.

سبحان اللّه و بعدد



Cells of the immune system / Lymphocytes/ T-lymphocytes


اللهم حلّ ملى يتد

• T lymphocytes, the cells of cell-mediated immunity, recognize the antigens of intracellular microbes and either help phagocytes to destroy these microbes or directly kill the infected cells.

(s) cytotoxicTcells

- T<u>cells do not produce antibo</u>dy من الميها تقتل صاطحة molecules.
- Their antigen receptors are membrane molecules distinct from but structurally related to antibodies.

سبحان الله و بعدد




microbes

of infection

Cell-mediated

### Cells of the immune system / Lymphocytes

- Humoral immunity is mediated by molecules in the blood and mucosal secretions, called antibodies, which are produced by cells called B lymphocytes.
- Cell-mediated immunity is mediated by T lymphocytes. viruses and some bacteria, survive and proliferate inside phagocytes and other host cells, where they are inaccessible to circulating antibodies. Defense against such infections is a function of cell mediated immunity, which promotes the destruction of microbes residing in phagocytes or the killing of infected cells to eliminate reservoirs of infection.



### Cells of the immune system / Lymphocytes/ T-lymphocytes

How do Teells cause cell apoptosis?

م يحد ثوا ثقوب في الخلايا المهافعة

Through the action of **perforin**. + **granzymes** enter the cytoplasm of the target cell and their serine protease function triggers the **caspase** cascade, which eventually lead to **apoptosis** (programmed cell death).

مرياه إلى الله والعليّ العظميم



#### Cells of the immune system / Lymphocytes





FIGURE 2–5 Maturation of lymphocytes. Lymphocytes develop from bone marrow stem cells and mature in the generative lymphoid organs (bone marrow and thymus for B and T cells, respectively) and then circulate through the blood to secondary lymphoid organs (lymph nodes, spleen, regional lymphoid tissues such as mucosa-associated lymphoid tissues). Fully mature T cells leave the thymus, but immature B cells leave the bone marrow and complete their maturation in secondary lymphoid organs. Naive lymphocytes may respond to foreign antigens in these secondary lymphoid tissues or return by lymphatic drainage to the blood and recirculate through other secondary lymphoid organs.

## Cells of the immune system / Lymphocytes


- The total number of lymphocytes in a healthy adults about  $5 \times 10^{11}$ . Of these:
  - $\sim$ 2% are in the blood,
  - $\sim$ 10% in the bone marrow,
  - ~15% in the mucosal lymphoid tissues of the gastrointestinal and respiratory tracts, and
  - $\sim$ 65% in lymphoid organs (mainly the lymph nodes and spleen)



**Further reading:** 

ر سنعنی سک العظیم و آخو ایس لایله الاست

• Cellular and Molecular Immunology. 7th Edition.. Chapter 2. Cells and tissues of the immune system



غيري لإخوا سي المسلمين ، أجلووا بقدر ما ستطيعون الدأوا عاندسكم وأمروا ما لعجون و الفو اعن الهنكر ولو كلقكم ذلاك الكيش