
سم الله الرحما الرحم

Medical Immunology



Anas Abu-Humaidan M.D. Ph.D. Page 2 of 27

### **Sepsis**



In this lecture we will discuss:

- When immunity becomes harmful
- The epidemiology and definition of sepsis
- The immunopathology of sepsis



## Introduction: When Immunity Becomes Harmful

- The immune system is essential for defense against pathogens.
- However, excessive, misdirected, or dysregulated responses can damage host tissues.

## 1. Molecular Mimicry and Autoimmune Diseases

- Pathogen antigens resemble self-proteins, leading to cross-reactivity.
- The immune system mistakenly attacks self-tissues.
- Examples:
  - Streptococcus pyogenes → rheumatic fever (cross-reactivity with heart tissue)
  - Campylobacter jejuni → Guillain-Barré syndrome (nerve glycolipid mimicry)
  - Coxsackie virus → Type 1 diabetes (molecular mimicry with islet antigens)

### 2. Autoimmune Mechanisms

- Breakdown of self-tolerance due to genetic, environmental, or infectious triggers.
- Mechanisms include:
  - Defective negative selection of autoreactive T/B cells.
  - Failure of regulatory T cells.
  - Epitope spreading during tissue damage.
- Examples: systemic lupus erythematosus, multiple sclerosis, rheumatoid arthritis.

## 3. Chronic Inflammation and Tissue Damage

- Persistent infection or autoimmunity leads to chronic immune activation.
- Continuous cytokine and ROS release damages tissues.
- Examples:
  - Chronic hepatitis → liver fibrosis.
  - Helicobacter pylori infection → gastritis, gastric cancer.
  - Tuberculosis → caseating granulomas causing lung damage.

# 4. Host Damage by Antiviral and Antitumor Immunity

- Cytotoxic T cells destroy infected or transformed cells.
- Collateral damage to nearby healthy tissues.
- Examples:
  - Viral hepatitis (HBV, HCV): immune-mediated hepatocyte injury.
  - Tumor immunity: inflammatory destruction of surrounding tissue.
- Balance between protection and pathology is critical.

Page 8 of 27

## 5. Cytokine Storms and Systemic Inflammation

- Overactivation of immune cells causes excessive cytokine release.
- Results in systemic inflammation, vascular leakage, and organ failure.
- Seen in severe viral infections (influenza, COVID-19, dengue) and bacterial sepsis.
- Key mediators: TNF- $\alpha$ , IL-1 $\beta$ , IL-6, IFN- $\gamma$ .
- Treatment: corticosteroids, cytokine inhibitors, supportive care.

## 6. Sepsis and Septic Shock

- Life-threatening organ dysfunction caused by dysregulated host response to infection.
- Triggered by microbial PAMPs activating PRRs on immune cells.
- Pathogenesis:
  - Massive cytokine release → vasodilation and capillary leak.
  - Coagulation activation  $\rightarrow$  DIC.
  - Mitochondrial dysfunction → organ failure.
- High mortality despite antimicrobial therapy.

#### Sepsis/overview

- Sepsis is a highly heterogeneous syndrome that is caused by an unbalanced host response to an infection.
- Sepsis was not clinically defined until the early 1990s when a group of key opinion leaders
  released the first consensus definition of sepsis. Since then, the definition was updated
  several times.
- The definition of sepsis matters in the **prognostication** of patients since those labelled as having sepsis are expected to have a **difficult clinical course and poor outcome** compared to those without sepsis.
- For many years, a disproportionate inflammatory response to invasive infection was considered to be central to the pathogenesis of sepsis, but it is now clear that the host response is disturbed in a much more complex way, involving both sustained excessive inflammation and immune suppression, and a failure to return to normal homeostasis

Page 11 of 27

### Sepsis/overview



991 Consensu	us Conference <sup>2</sup>
Diagnosis	Signs and symptoms
Systemic inflammatory response syndrome	Patients experiencing at least two of the following symptoms:  **Body temperature >38°C or <36°C  **Heart rate >90 beats per minute  **Respiratory rate >20 breaths per minute or arterial CO <sub>2</sub> <32 mmHg  **White blood cell count >12 × 10°l' ar <4 × 10°l' ar, or >10%  **Immature forms  **Least W. J. L. W.
Sepsis	Systemic inflammatory response syndrome and proven or suspected infection
Severe sepsis	Sepsis and acute organ dysfunction
Septic shock	Sepsis and persistent hypotension after fluid resuscitation

2001 International Sepsis Definitions Conference<sup>145</sup>
The 2001 definitions of sepsis were very similar to the definitions stated in 1991.
Of note, in 2001 it was acknowledged that the signs and symptoms of sepsis are more varied than described in the 1991 definition, and this resulted in the addition of a list of these signs and symptoms for the diagnosis of sepsis.

Diagnosis	اللقى بين الحد يدوال عم يا Signs		
Sepsis*	<ul> <li>Life-threatening organ dysfunction caused by a dysregulated host response to infection</li> <li>Organ dysfunction can be identified as an acute change in total SOFA score of ≥2 points<sup>‡</sup></li> </ul>		
Septic shock	<ul> <li>Sepsis in which the underlying circulatory and cellular and/or metabolic abnormalities are marked enough to substantially increase mortality</li> <li>Clinically defined as sepsis with persisting hypotension that require vasopressors to maintain the mean arterial pressure at ≥65 mmHg and with a serum lactate concentration &gt;2 mmol</li> </ul>		

\*Of note, the presence of organ dysfunction is central and required in the new 2016 consensus sepsis definition. Until then, organ dysfunction was part of the definition of 'severe' sepsis, a term that was abandoned in the Sepsis-3 definition. \*The sequential organ failure assessment (SOFA) score is based on six different scores (each classified from 1 to 4 according to increasing abnormality and/or severity), one each for the respiratory, cardiovascular, hepatic, coagulation, renal and neurological systems146.

Page 12 of 27

### Sepsis/overview

## اللهم مِلَّ وسلِّم على نبيّنا في ل

### مش مهم نعرض س مراء ه

SOFA score	0	1	2	3	4
Respiration	2.2 (0.00)	Outdoor (Observation			S. 400
PaO <sub>2</sub> /FIO <sub>2</sub> (mmHg)	> 400	301-400	201-300	101-200	≤ 100
(kPa)	> 5.3)	(4.1-5.3)	(2.8-4.0)	(1.4-2.7)	≤ 1.3)
Coagulation					
Platelets (x10 <sup>3</sup> /mm <sup>3</sup> )	> 150	101-150	51-100	21-50	≤ 20
Liver					
Bilirubin (mg/dl)	< 1.2	1.2-1.9	2.0-5.9	6.0-11.9	≥ 12.0
(µmol/l)	< 20)	(20-32)	(33-101)	(102-204)	≥ 204)
Cardiovascular					
Hypotension	No hypotension	MAP < 70 mmHg	Dopamine ≤ 5 or dobutamine (any dose)*	Dopamine > 5	Dopamine > 15
Central nervous system					
Glasgow coma score	15	13-14	10-12	6-9	< 6
Renal					
Creatinine (mg/dl)	< 1.2	1.2-1.9	2.0-3.4	3.5-4.9	> 5.0
(µmol/l)	< 110)	(110-170)	(171-299)	(300-440)	> 440)
or urine output	THE PERSON NAMED IN COLUMN 1	0.000 0.00000 0.0000000	Marie Control Control No.	< 500 ml/day	< 200 ml/day

<sup>\*</sup> adrenergic agents administered for at least 1 h (doses given are in µg/kg/min)

### Sepsis/epidemiology

لاحول ول حَوِّج إلَّه عالله

- For 2017, it was estimated that it had affected 49 million individuals and was related to approximately 11 million potentially avoidable deaths worldwide.
- Sepsis mortality is often related to suboptimal quality of care, an inadequate health infrastructure, poor infection prevention measures in place, late diagnosis, and inappropriate clinical management.
- Antimicrobial resistance further complicates sepsis management across all settings, particularly in high-risk populations, such as neonates and patients in intensive care units (ICUs)
- One in four cases of sepsis in hospitals and one in two cases of sepsis in ICUs result from health care-associated infections.

### Sepsis/epidemiology

## Characteristics of Adult Sepsis Patients in the Intensive Care Units in a Tertiary Hospital in Jordan: An Observational Study

**Anas H. A. Abu-Humaidan**  $\bigcirc$   $\bigcirc$ , <sup>1</sup> Fatima M. Ahmad  $\bigcirc$ , <sup>1,2</sup> Maysaa' A. Al-Binni, <sup>2</sup> Amjad Bani Hani  $\bigcirc$ , <sup>3</sup> and Mahmoud Abu Abeeleh  $\bigcirc$ 3

All adult patients admitted to the adult ICUs between June 2020 and January 2021 were included in the study. Patients' clinical and demographic data, comorbidities, ICU length of stay (LOS), medical interventions, microbiological findings, and mortality rate were studied.

We observed 194 ICU patients during the study period; 45 patients (23.3%) were diagnosed with sepsis using the Sepsis-3 criteria.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8736695/



Mortality rate and median ICU LOS in patients who had sepsis were significantly higher than those in other ICU patients (mortality rate, 57.8% vs. 6.0%, value < 0.001, resp., and LOS 7 days vs. 4 days, value < 0.001, resp.).

Additionally, sepsis patients had a higher combined number of comorbidities. The use of mechanical ventilation, endotracheal intubation, and blood transfusions were all significantly more common among sepsis patients.

Microbiological findings in sepsis patients.

<sup>1</sup> Suspected origin of infection, n (%) Gastrointestinal, 17 (37.8%)

Respiratory, 11 (24.4%)

Genitourinary, 11 (24.4%)

Skin and soft tissue, 6 (13.3%)

Others, 16 (17.8%)

<sup>2</sup> Isolated organisms, n (%)

Gram-positive bacteria, 17 (35.4%)

Staphylococci (coagulase-negative), 10 (20.8%)

Staphylococci (coagulase-positive), 2 (4.2%)

 $\it Enterococcus \, species, 4 \, (8.3\%)$ 

Streptococcus species, 1 (2.1%)

Gram-negative bacteria, 37 (77.1%)

Escherichia coli, 10 (20.8%)

Acinetobacter baumani, 10 (20.8%)

Klebsiella species, 9 (18.8%)

Pseudomonas aeruginosa, 3 (6.3%)

Others, 5 (10.4%)

Fungi, 10 (20.8%)

Candida species, 10 (20.8%)

لاحول ول حَوِّج إلَّه عاللَّ

### Sepsis/The initiation of inflammation



- Sepsis is associated with a strong activation of the **innate immune system** that is mediated by the activation **of PRRs** by **PAMPs** and **DAMPs**
- There is similarity between the inflammatory reactions induced by different pathogens and those elicited by different types of injury, either infectious or non-infectious.
- Pro-inflammatory cytokines implicated in sepsis pathogenesis include **tumour necrosis** factor (TNF), interleukin-1 $\beta$  (IL-1 $\beta$ ), IL-12 and IL-18; blocking or eliminating these cytokines confers protection in acute animal models of fulminant infection.

### Sepsis/Complement activation



- Although complement activation is an essential component of protective immunity, the uncontrolled activation of complement can cause damage to tissues and organ failure.
- Activation of the three major pathways of complement, including the classical, lectin, and alternative pathways can take place in sepsis.
- Moreover, some studies reported a correlation between levels of complement activation fragments such as C3b and C5a to sepsis severity. Complement activation can culminate in activation of the terminal pathway, through cleavage of C5 to form C5a, a potent anaphylatoxin, and C5b, which initiates formation of the pore forming complex C5b-9 on cellular membranes.
- Blockade of C5a signalling improved the outcome of experimental sepsis in several animal models, including Escherichia coli sepsis in baboons and rats with polymicrobial abdominal sepsis

Page 18 of 27

#### Sepsis/Complement activation

الحد الله

Journal of Inflammation Research





ORIGINAL RESEARCH

### Complement Terminal Pathway Activation is Associated with Organ Failure in Sepsis Patients

Fatima M Ahmad (6).2 Maysaa' A Al-Binni<sup>2</sup> Amjad Bani Hani (6)<sup>3</sup> Mahmoud Abu Abeeleh<sup>3</sup> Anas HA Abu-Humaidan

<sup>1</sup>Department of Pathology, Microbiology and Forensic Medicine, School of Medicine, The University of Jordan, Amman, Jordan; <sup>2</sup>Department of the Clinical Laboratory Sciences, School of Science, The University of Jordan, Amman, Jordan; <sup>3</sup>Department of General Surgery, School of Medicine, The University of Jordan, Amman, Jordan

Conclusion: In sepsis patients, levels of C5 and sCD59, but not sC5b-9, correlated to the severity of organ damage measured by SOFA A similar correlation was not found in non sepsis patients. This indicated that organ damage associated with sepsis led to a more pronounced terminal pathway activation than in non-sepsis patients, it also indicated the potential of using C5 and sCD59 to reflect sepsis severity.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8760944/

Page 19 of 27



### Sepsis/Coagulation, endothelial cell activation and vascular leakage

- Sepsis is associated with a **strong activation of the coagulation system**, and this can result in disseminated intravascular coagulation, which clinically can be associated with microvascular thrombosis and haemorrhage, the latter being due to the consumption of clotting factors and platelets. **Tissue factor** is the main driver of coagulation activation in sepsis. In addition, tissue factor inhibition prevents multiple organ failure and mortality in a model of other- wise lethal sepsis in baboons.
  - In response to localized infection, leukocytes and platelets adhere to the endothelial surface and migrate to the sites at which bacteria are multiplying. In sepsis, exaggerated inflammation augments these processes, thereby con-tributing to barrier incompetency
  - A loss of barrier integrity causes the **leakage of intravascular proteins and plasma into the extravascular space**, tissue oedema and reduced microvascular perfusion.

4 then hypotension

### Sepsis/Other immune mechanisms



- **Neutrophil extracellular traps.** NETs can entrap pathogens and thereby contribute to pathogen elimination. However, NETs can also contribute to collateral tissue damage and thrombosis. Patients with sepsis have **increased NET levels** in their circulation, and this feature is associated with organ dysfunction.
- The role of platelets. Excessive platelet activation has been implicated in organ injury
  during sepsis through several mechanisms, including the augmentation of immune cell
  recruitment and inflammation, the facilitation of the formation of vaso-occlusive thrombi in
  capillary vascular beds and direct cell toxic effects mediated by platelet derived
  microparticles.
- The role of B cells. A subset of B cells, the so-called **innate response activator B cells**, is important for bacterial eradication as well as for the attenuation of proinflammatory cytokine release. Innate response activator B cells can produce IL-3, which in the context of sepsis increases inflammation and the production of myeloid mononuclear cells.

Sepsis/ Immune suppression in sepsis



- Sepsis is associated with immune suppression that is characterized by lymphocyte exhaustion and the reprogramming of antigen-presenting cells.
- Sepsis is associated with a strong **depletion** of CD4+ and CD8+ T cells, B cells and dendritic cells (DCs) as a result of apoptosis
- Immune suppression in sepsis is characterized by the **reduced expression of HLA-DR** on blood monocytes, and by the diminished capacity of monocytes and macrophages to release pro-inflammatory cytokines upon stimulation.
- The immune suppression increases the risk of secondary infections. A recent observational study indicated that secondary infections are responsible for only 10.9% of overall sepsis mortality in the ICU.


### Sepsis/ Overview


## اللهم علة وسلّم على نبيّنا محيّد

#### Protective immunity Localized innate immune • Rel<u>ease of</u> p<u>ro-inflam</u>matory • Release of pro-inflammatory Release of pro-inflammatory mediators mediators mediators • Leukocyte recruitment ↑ Adhesive and procoagulant properties ↓ Barrier function Cell injury with release of DAMPs Pro-inflammatory Complement activation Coagulation activation response PAMPs Coagulation activation (microvascular thrombosis) · Release of pro-inflammatory CLR mediators • Activation of neutrophils Complement activation and the endothelium • Microvascular thrombi TLRs **SNLR** Homeostasis --RLR CD4+ T cells CD8+ T cells • ↑ Apoptosis • ↑ Apoptosis Exhaustion ↓ Cytoxic function Exhaustion • T<sub>H</sub>2 cell polarization Local repair mechanisms Inhibition and resolution Anti-inflammatory mechanisms of inflammation ↓ Apoptosis Reprogramming of Tissue repair Return to homeostasis ↑ Immature cells with macrophages to an M2 decreased antimicrobial phenotype • Reduced HLA-DR functions expression

Apoptosis of B cells and follicular DCs

 Expansion of regulatory T cell and MDSC populations





Pathological alteration and therapeutic implications of sepsisinduced immune cell apoptosis

### Sepsis/ Immunomodulation as a treatment for sepsis



- How the host response should be manipulated in patients with sepsis is controversial. The immune disturbances are complex and require targeting more than one pathway/ mechanism.
- Immune suppression through inhibition of complement or coagulation (examples, C5a-specific monoclonal antibody, recombinant human thrombomodulin)
- Blood purification techniques have been proposed as a method of removing PAMPs and inflammatory mediators from the circulation of patients. Recently, a blood-cleansing device that removes multiple pathogens and toxins from the blood via magnetic nanobeads coated with an engineered form of the human opsonin mannose-binding lectin (also known as MBPC) was described, and it is currently being evaluated in preclinical studies
- Immune stimulation. There are several drugs that could potentially reverse immune suppression in sepsis. One approach is to use immune-stimulating cytokines, such as IFNγ, IL-7 and IL-15.

# Summary Table: Detrimental Immune Mechanisms

Mechanism	Example Disease	Main Mediator	Effect
Molecular mimicry	Rheumatic fever	Cross-reactive antibodies	Myocardial injury
Cytokine storm	Sepsis, COVID-19	TNF-α, IL-6	Multiorgan failure
Immune complex deposition	SLE, GN	IgG, complement	Vasculitis, nephritis
Chronic inflammation	TB, H. pylori	Macrophages, TNF-α	Fibrosis
CTL-mediated damage	Viral hepatitis	CD8+ T cells	Hepatocyte death

### Key Takeaways



- Immune responses can inadvertently harm the host through autoimmunity, hyperinflammation, or chronic activation.
- Understanding immune-mediated pathology is essential for designing safer vaccines and therapies.
- Regulation, not suppression, is key to maintaining immune balance.

### **Further reading:**

أستنعر الله العظم وأكون إليه

• The immunopathology of sepsis and potential therapeutic targets. *Nature Reviews Immunology volume 17, pages407–420 (2017)* 



کی آجتم بالکومیت به انظیمی ، ساء علی عظم أوحفظ أوسالم وقرادة آیانا من الفرآن سكل بودي ٥



التوفيق، ادمولاا.