
and the West Werd

Medical Immunology



Anas Abu-Humaidan M.D. Ph.D.

Lecture 7

Antibodies and antigens

In this lecture we will discuss:

• Antibodies, their structure, types and function.

اللهم مِل وسلّم على سينا محسد


Page 3 of 30

Antibodies/ overview

لل حول دلاحرة إلَّد بالله العطيم

all serum profeins


- The study of antibodies and their reactions with antigens is therefore classically called serology.
- **Serum** lacks coagulation factors but otherwise contains all the proteins found in plasma. Any serum sample that contains detectable antibody molecules that bind to a particular antigen is commonly called an **antiserum**.
- A healthy 70-kg adult human produces about 2 to 3 g of antibodies every day.



Page 4 of 30

Antibodies/ overview


• Plasma proteomics holds great promise for the future of biomarker discovery, as well as *in vitro* diagnostics. Although plasma is readily accessible for analysis, the study of the plasma proteome is fundamentally limited by its vast dynamic range (10 orders of magnitude).



Antibodies and antigens

- Antibodies are circulating proteins that are produced in vertebrates in response to exposure to foreign structures known as antigens
- The strength of the binding between a single site of an antibody and an epitope of an antigen is called the affinity of the antibody.

لا حول دلاتية إلَّا الله العطيم




Antibodies and antigens

• Because the hinge region of antibodies gives them flexibility, a single antibody may attach to a single multivalent antigen by more than one binding site. the strength of attachment of the antibody to the antigen must take into account binding of all the sites to all the available epitopes. This overall strength of attachment is called the avidity.


Posted poi affinity)


اللهم مل على بينا عدد




Antibodies/structure

- Antibodies (immunoglobulins) are made of 2 heavy chains and 2 light chains, those chains combined give us an antibody binding region (Fab) and the fragment crystallizable region (Fc region) which is the tail region of an antibody that interacts with cell surface receptors called Fc receptors and some proteins of the complement system.
 - Immunoglobulins are divided into different classes (isotypes).








Antibodies/ structure

اللهم مهل وسلم على بنينا في ر

- Antibodies can exist in two forms:
 membrane-bound antibodies on the
 surface of B lymphocytes function as
 receptors for antigen, and secreted
 antibodies that reside in the circulation,
 tissues, and mucosal sites
- Both heavy chains and light chains consist of amino terminal variable (V) regions that participate in antigen recognition and carboxyl-terminal constant (C) regions; the C regions of the heavy chains mediate effector functions.



Remember Igs are proteins so these are

poly peptide chains (2 light + 2 Heavy)

للهاله إلا الله


- Question: How many polypeptide chains make up a typical antibody molecule?
- a) 1
- b) 2
- c) 3
- d) 4

Page 10 of 30

Antibodies/ structure

لا حول دلاتية إلَّد الله العطيم

Most of the sequence differences and variability among different antibodies are confined to three short stretches in the V region of the heavy chain and to three stretches in the V region of the light chain. These diverse stretches are known as hypervariable segments, Because these sequences form a surface that is complementary to the three-dimensional structure of the bound antigen, the hypervariable regions are also called complementarity-determining regions (CDRs).




Page 11 of 30

Antibodies

للايله إلا الله

- There are five immunoglobulin classes (isotypes) of antibody molecules found in serum: IgG, IgM,
 IgA, IgE and IgD. They are distinguished by the
 type of heavy chain they contain
- Antibodies of different classes differ in their location around the body, appear at different stages of an adaptive immune response.


• The heavy chain C regions of all antibody molecules of one isotype or subtype have essentially the same amino acid sequence. This sequence is different in antibodies of other isotypes or subtypes.



Antibodies/ structure

	sotope of Antibody	Subtypes (H Chain)	Serum Concentration (mg/mL)	Serum Half-life (days)	Secreted Form		Functions
I.	gA	IgA1,2 (α1 or α2)	3.5	6	IgA (dimer) Monomerkdimer, trimer the most Common Form	Cut	Mucosal immunity (3 in mucosal frac
- In	gD	None (δ)	Trace	3	None		Naive B cell antigen receptor
l,	gE	None (ε)	0.05	2	IgE Monomer	CD Co2 CD Co2 CD Co3 CD Co4	Defense against helminthic parasites, immediate hypersensitivity
3 4	gG	1gG1-4 (γ1, γ2, γ3, or γ4) Camma 4	13.5 1xpes	23	lgG1 Monomer	45.50 % A STATE OF THE PARTY OF	Opsonization, complement activation, antibody- dependent cell-mediated cytotoxicity, neonatal immunity, feedback inhibition of B cells
(J)	gM	None (μ)	1.5	5	IgM Pentamer	CH1, CH2	Naive B cell antigen receptor, complement activation

لا حول ولاحرة إلَّد بالله العطيم



• Question: Which is the most abundant Ig class in serum?


- a) IgG
- b)IgA
- c) IgM
- d) IgE


B CELLS AND ANTIBODY RESPONSES

B lymphocytes are the cells responsible for antibody responses.

genefic recombination


- During its development, each B-lymphocyte becomes genetically programmed through a series of gene-splicing reactions to produce an antibody molecule with a unique specificity
- It is estimated that the human body has the ability to recognize 10⁷ or more different epitopes, due to the wide range of possible combinations during gene splicing.





Page 15 of 30


B Cell Activation



- Resting B cells become activated by antigen via binding to the BCR and internalization.
- Once internalized inside the B cell, the protein antigen is processed and presented with MHC II. The presented antigen is then recognized by helper T cells specific to the same antigen.
- Once activated by linked recognition, **T-cells produce and secrete cytokines that activate the B cell and cause proliferation** into clonal daughter cells.
- After several rounds of proliferation, additional cytokines provided by the T-cells stimulate
 the differentiation of activated B cell clones into memory B cells, which will quickly
 respond to subsequent exposures to the same antigen, and plasma cells that lose their
 membrane BCRs and initially secrete pentameric IgM

Page 16 of 30

B Cell Activation




Antibodies/ Antibody Production






Initial contact with a new antigen evokes the **primary response**, which is characterized by a lag phase of approximately 1 week between the challenge and the detection of circulating antibodies.

- Once antibody is detected in serum, the levels rise exponentially to attain a maximal steady state in approximately 3 weeks, then decline gradually with time.
- The **first antibodies** synthesized in the primary immune response **are IgM** and, **then IgG** antibodies arise and eventually predominate.



Antibodies/ Antibody Production




2

 After a subsequent exposure or booster injection of the same antigen, a different sequence called the secondary response ensues.

• In the secondary response; العزرهات

the **lag time** between the immunization and the appearance of antibody is **shortened**,
- the rate of exponential increase to the maximum steady-state level is **more rapid**,
and the steady-state level itself is higher, representing a **larger amount of antibody**.

 Another key factor of the secondary response is that the antibodies formed are predominantly of the IgG class. Page 19 of 30



سعان الله و بعدم سحان (لله العطيم


Antibodies/ Class switching

حسبن الله ونحم الوكيل

- After initial secretion of IgM, cytokines secreted by T- cells stimulate the plasma cells to switch from IgM production to production of IgG, IgA, or IgE.
- This process, called class switching or isotype switching, allows plasma cells cloned from the same activated B cell to produce a variety of antibody classes with the same antigen specificity.
- Class switching is accomplished by genetic rearrangement of gene segments encoding the constant region, which determines an antibody's class. The variable region is not changed, so the new class of antibody retains the original antigen specificity.


Class switch

- $\ensuremath{\mathbf{DNA}}$ rearrangement changing the heavy chain constant gene in memory cells.




Page 21 of 30

Antibodies and antigens




- The ability of antibodies in any individual to specifically bind a large number of different antigens is a reflection of antibody diversity, and the total collection of antibodies with different specificities represents the antibody repertoire. The genetic mechanisms that generate such a large antibody repertoire occur exclusively in lymphocytes. This diversity is generated by random recombination of a limited set of inherited germline DNA sequences to form functional genes that encode the V regions of heavy and light chains as well as by the addition of nucleotide sequences during the recombination process.
- Somatic mutation in antigen-stimulated B lymphocytes that generates new V domain structures, some of which bind the antigen with greater affinity than did the original V domains. Those B cells producing higher affinity antibodies preferentially bind to the antigen and, as a result of selection, become the dominant B cells with each subsequent exposure to the antigen. This process is called affinity maturation.



Page 23 of 30

Effector mechanisms of humoral immunity

Neutralization & ABC




أُستَعَفَى اللَّهُ العظيم عدد ما حلق و قل عا حلق

Antibodies/ IgM

- Because of its many specific binding sites, IgM is particularly effective in agglutinating particles carrying antigens against which it is directed.
- IgM is particularly active in bringing about complement-mediated cytolytic damage to foreign antigen-bearing cells.


لله إلى إلى الله وحده لا شريك له له الله الله وهو على كلّ شي (م) به



Antibodies/ IgG

أُستَعَفَراللهُ والعظيم عدد ما حلق و قل والحلق

- Immunoglobulin G (IgG) is the most abundant immunoglobulin in health and provides the most extensive and long-lived antibody response to the various microbial and other antigens.
- IgG antibody is characteristically formed in large amounts during the secondary response to an antigenic stimulus, and usually follows production of IgM in the course of a viral or bacterial infection.
- IgG is the only immunoglobulin class able to cross the placental barrier and, thus, provides passive immune protection to the newborn in the form of maternal antibody.




Antibodies/ IgA


سسان الله و بعده سحان الله العطيم

- Immunoglobulin A (IgA) has a special role as a major determinant of so-called local immunity in protecting epithelial surfaces from colonization and infection.
- At the epithelia, two IgA molecules combine with another protein, termed the secretory piece, which is present on the surface of local epithelial cells. The complex, then termed secretory IgA (sIgA),

of is reallered on is Early

 The major role of slgA is to prevent attachment of antigen-carrying particles to receptors on mucous membrane epithelia








Antibodies/ IgE حلل مَن وَ وَلَا اللهِ عَلَى اللهِ عَلَى اللهِ عَلَى اللهِ عَلَى اللهِ عَلَى اللهِ عَلَى اللهِ


IgE not only provides protective immunity
against helminth parasites but can also mediate
the type I hypersensitivity reactions that
contribute to the pathogenesis of allergic
diseases such as asthma, allergic rhinitis and
atopic dermatitis.



Antibodies/ IgD

- IgD (IgD) is a monomeric antibody isotype that is expressed in the plasma membranes of immature B-lymphocytes. IgD is also produced in a secreted form that is found in small amounts in blood serum.
- the function of IgD is to signal the B cells to be activated.

أُستَعَفَى اللَّهُ العظيم عدد ما حلق و قل عا حلق



. It goes class switching to Igh, IgG, IgF

Page 29 of 30



 $\frac{https://www.jacksonimmuno.com/secondary-antibody-resource/technical-tips/immunoglossary/\#:^:text=Complementarity%20Determining%20Regions%20(CDRs)&text=An%20lgG%2Ontibody%20has%2012,each%20of%20the%20heavy%20chains.$

Further reading:

Cellular and Molecular Immunology. 7th Edition..
 Chapter 5. Antibodies and antigens

للفائدةولعلّها تكون سببًا في تعجيل النّصر أو إزاحة الهموم عن إخواننا بإذن الله: كان النبي صلى الله عليه وسلم يدعو في وقت الأزمات والابتلاءات التي تصيب المسلمين بعد القيام من الركوع الأخير في كلّ صلاة مفروضة وهذا يسمّى القنوت ويستحب فعله في المساجد فيدعو الإمام ويُؤمّم المصلّون .**الدعاء يكون في المساب فيدعو للمسلمين المستضعفين في المصاب فيدعو للمسلمين المستضعفين ويدعو على الظالمين ولا تشترط الإطالة في الدّعاء الدّعاء الدّعاء الدّعاء