




HYPERTENSIVE VASCULAR DISEASE

Arteriolosclerosis

Dr. Nisreen Abu Shahin
Professor of Pathology
Pathology Department
University of Jordan



A sphygmomanometer or a Digital blood pressure monitor is used to measure BP.

Types of hypertension

- **According to severity:**

Benign (95%) versus malignant (5%)

- **According to cause:**

Primary (essential) (95%) versus secondary (5%)

- **Another way to classify:**

Systolic vs diastolic

- ***Malignant hypertension***

- 5% (also known as accelerated HTN)

- a rapidly rising blood pressure that, if untreated, leads to death within 1 to 2 years


- **systolic pressures > 200 mm Hg or diastolic pressures > 120 mm Hg**

- renal failure and retinal hemorrhages

- usually superimposed on preexisting **benign** hypertension (either essential or secondary)

Hypertension (HTN) has the following potential complications:

Complications of Hypertension: Target-Organ Damage



TIA, transient ischemic attack; LVH, left ventricular hypertrophy; CHD, coronary heart disease; HF, heart failure



- stroke (CVD) & multi-infarct dementia
- atherosclerotic coronary heart disease
- cardiac hypertrophy and heart failure (*hypertensive heart disease*)
- aortic dissection
- renal failure
- retinal hemorrhages

Types of hypertension- according to etiology

- 1- **essential (idiopathic) hypertension (95%)**
- 2- **secondary hypertension:**
 - **Most common of secondary:** renal disease or renal artery narrowing (**renovascular hypertension**)
 - **Other less common:** many other conditions....

Essential Hypertension

Accounts for 90% to 95% of all cases

Most common of all

Secondary Hypertension

Renal

Acute glomerulonephritis

Chronic renal disease

Polycystic disease

Renal artery stenosis

Renal vasculitis

Renin-producing tumors

Most common of
secondary causes

Endocrine

Adrenocortical hyperfunction (Cushing syndrome, primary aldosteronism, congenital adrenal hyperplasia, licorice ingestion)

Exogenous hormones (glucocorticoids, estrogen [including pregnancy-induced and oral contraceptives], sympathomimetics and tyramine-containing foods, monoamine oxidase inhibitors)

Pheochromocytoma

Acromegaly

Hypothyroidism (myxedema)

Hyperthyroidism (thyrotoxicosis)

Pregnancy-induced (pre-eclampsia)

Cardiovascular

Coarctation of aorta

Polyarteritis nodosa

Increased intravascular volume

Increased cardiac output

Rigidity of the aorta

Neurologic

Psychogenic

Increased intracranial pressure

Sleep apnea

Acute stress, including surgery



- **Pathogenesis of essential HTN**

- **? Genetic factors**

? familial clustering of hypertension

- angiotensinogen **polymorphisms** and angiotensin II receptor variants; polymorphisms of the renin-angiotensin system.
- ? **Susceptibility** genes for essential hypertension: genes that control renal sodium absorption, etc...


- **Environmental factors** modify the impact of genetic determinants

stress, obesity, smoking, physical inactivity, ↑ salt consumption

Blood vessels in HTN- Morphology

- HTN is associated with **arteriolosclerosis** (small arterial disease)
- Two forms of small blood vessel disease are hypertension-related:
 - 1- **hyaline arteriolosclerosis**
 - 2- **hyperplastic arteriolosclerosis**

1- Hyaline arteriolosclerosis



- Ass. with benign hypertension
- homogeneous **pink** hyaline thickening of arteriolar walls
- luminal narrowing
- leakage of plasma components across injured endothelial cells into vessel walls
- increased ECM production by smooth muscle cells in response to chronic hemodynamic stress

- **Hyaline arteriolosclerosis: Complications**


- **Most significant in kidneys → nephrosclerosis (glomerular scarring)**

- Other causes of **hyaline** arteriolosclerosis:

1- elderly patients (normo-tensive)


2- diabetes mellitus

2- Hyperplastic arteriolosclerosis



- With severe (malignant) hypertension
- "onionskin" concentric laminated **thickening** of arteriolar walls
- luminal narrowing
- **reduplicated** basement membrane
- fibrinoid vessel wall necrosis (necrotizing arteriolitis)

Fibrinoid Necrosis - artery



Q: What does (reduplicated basement membrane) mean ?

A: It means that the basement membrane will have multiple layers instead of the normal one, so it becomes thicker than normal and is functionally considered abnormal and results in reduced blood flow to the tissue.

The presence of these multiple layers will give the appearance of the onion when it is cut. This is termed (onion skin) appearance.