




HISTOLOGY

LEC : 1+2


LEC1 : Histology of upper respiratory tract.

- ☆ Respiratory System is made of ① Conducting ② Respiratory parts.
- Conducting: Nose + pharynx + larynx + trachea + Rt. and Lt. main bronchi (primary) + secondary bronchi (Lobar bronchi: Branch for each lobe → 3 in Right and 2 in left lung).
- + 10 bronchopulmonary segments on each side (tertiary bronchi).
- ↳ Each segment is surrounded by connective tissue and contains a bronchi + blood vessels lymphatics.

Zones of the Respiratory System



After Bronchi → Bronchioles: 1- Very narrow diameter



2- Different type of epithelium

3- No cartilage

4- Contain spiral helical smooth muscles.

5- Classified to $\left\{ \begin{array}{l} \text{large conducting bronchioles} \\ \text{terminal bronchioles} \end{array} \right.$

Terminal bronchioles: Immediately before respiratory portion.

↳ ① No cartilage ② No glands ③ No goblet cells

↳ Epithelium: Simple cuboidal with or without cilia (Clara cells).

↳ Simple cuboidal without goblet cells.

Respiratory portion: Simple squamous epithelium

↳ Starts as respiratory bronchioles that's narrower than terminal bronchioles.

- Respiratory bronchioles open in alveolar ducts that's surrounded by alveoli.
- Alveolar duct terminate as Alveolar sac.
- Alveoli surround the ① Respiratory bronchioles ② Alveolar ducts ③ Alveolar sacs.
- Each alveolus is surrounded by a network of capillaries.

- In the beginning of the Respiratory portion there will be some Clara cells.

Conducting portion

layer 1: Mucosa

Mucosa → lining epithelium
→ lamina propria (No glands or very minimal)
→ Muscularis mucosa

Functions: ① Conducts
② warms ③ moistures
④ cleans the air

☆ Lining epithelium: Pseudostratified Columnar Ciliated with goblet cells

→ From nose → toward tertiary bronchioles.

→ Simple Cuboidal with or without goblet cells
→ terminal bronchioles.

layer 2: Submucosa

Contain Seromucous gland with a duct opens in the lumen

layer 3: Subcartilage layer

☆ Hyaline Cartilage and Smooth muscles.

→ For keeping the pathway open for easy passage of air.

In trachea: C-shaped hyaline cartilage. Absent posteriorly → trachealis

smooth muscles posteriorly. why?

Because posterior to the trachea there is the esophagus: if there were cartilage it may stop the passage of bolus in esophagus.

☆ In primary + secondary + tertiary bronchi: plates of cartilage.

- Decrease when going distally.

In 1° it's surrounding the lumen in all directions.

In 2° it's less present

In 3° you may only find 1 or 2 plates

While Smooth muscles is fewer in areas with higher content of cartilage like trachea

- In tertiary bronchi: there will be a lot of smooth muscles.

- In bronchioles it's heavy, spiral and helical surrounding all the lumen.

Layer 4: Adventitia (Connective tissue)

Respiratory portion

* Alveoli: The wall is made of 2 types of cells

☆ Type 1: 97%
→ Important for gas exchange.

☆ Type 2: 3% (Cuboidal cells).
→ Function: Secretion of a protein known as **Surfactant** → decrease surface tension in alveolus → easier inflation.

Before birth: the levels of surfactant must be measured.

☆ When the baby is born the doctor gently taps baby's back to stimulate phrenic nerve → Impulses go to respiratory centers in the brain → Motor impulses through phrenic nerve to diaphragm → diaphragm descends downward → lower intrapleural pressure → inspiration of air.



- Surfactant helps lungs to be filled with air, if it was insufficient → baby won't breathe Infant Respiratory Distress Syndrome (IRDS)

Respiratory Membrane


☆ Blood-Air Barrier

☆ Composed of :

- ① Surfactant
- ② Endothelial cell wall
- ③ Type 1 Alveolar Cell wall
- ④ Fused basal lamina (from blood capillary and Alveolus)



Respiratory Epithelium



Pseudostratified ciliated columnar Epithelium with Goblet cells

- Seen under light microscope

- 300 cilia on its apical surface.

↳ moves in one direction (outward)



↳ Attached to basal body

Goblet cells
secrete mucus.
for moistening
and filtration

- under the basal bodies there are Apical Mitochondria.

↳ Abundant in large numbers

↳ gives energy to cilia for motility



Ciliary Movement

Dynin : protein participate in the ciliary movement.
located near the basal bodies.

☆ Smoking leads to the release of high amounts of nicotine that leads to destruction of Dynein or inhibits its synthesis → destroyed ciliary movement (Immotile cilia syndrom (Kartagener Syndrom)).

Any pathogen enters will persist in the respiratory tract leading to chronic infections.

In males, dynein also helps in sperms motility. Smoking destruct sperms motility → Infertility.

* under Electron microscope we're able to see other types of cells.

① Basal cells : on the basement membrane and don't reach the surface.



Reserve cell → high mitotic activity.

② Brush cells: - Contain microvillie on apical surface
- Sensory receptors

③ Serous cells: (Granular) Contain secretory granules.

④ Diffuse neuroendocrine cells. Kulchitsky cells: Regulate secretion of glands

⑤ Goblet cells.



Nasal Cavity


Subdivided into:

① Vestibule

Epithelium: Stratified squamous keratinized (like skin)

Contain vibrissae: thick hair for filtration

Contain sebaceous glands



② Respiratory area

- Below Superior Concha and above the vestibule.

- Epithelium: pseudostratified ciliated columnar with goblet cells.

- Submucosa contain plexus of veins.

③ Olfactory region

- Above the Superior Concha (Roof).

- Epithelium: pseudostratified ciliated columnar with goblet cells

- Contain bipolar cells: one pole contain hairlike


The other pole have filament of olfactory nerve.

Converts odor into smell impulses → Smell centers

- In the olfactory region Glands are both in lamina propria and submucosa.

- The glands have ducts toward the mucus surface.
These Glands are Bowman's Gland

- Contain Supporting (Sustentacular) columnar cells
- Contain Basal cells



Nasal Sinuses

- lined by thin mucus that's adherent with the pericostum.
- Pseudostratified columnar with few goblet cells.
- and few glands

Structural Changes in the Bronchial Tree









- ① Goblet cells: Decrease when going distally then become absent.
- ② Glands: Decrease when going distally then become absent.
- ③ Smooth muscles: Increase when going distally (most prominent in Bronchioles) and in lungs become as patches and replaced by Elastic and Reticular fibers.
- ④ Elastic and Reticular fibers: present every where but most prominent
(In lungs between alveoli).
↳ Responsible for inflation and deflation.
- ⑤ Lymphocytes: Scattered everywhere and aggregates and form lymphatic Nodule around bronchioles.

Cough Reflex

Any foreign body entering the respiratory tract will induce cough to push it outside

Extra-pulmonary Bronchi:
Are the primary bronchi. (outside the lungs).

Intra pulmonary Bronchi:
Are the secondary and Tertiary (Surrounded by lung tissue).



Nasal Sinuses

LEC-2

Lower Respiratory Tract

Bronchioles


- Diameter 5mm → then divides more → large bronchioles are 1mm.
→ Small are 0.5mm.
- No cartilage + No glands + Few Goblet cells.

→ Clara cells (Simple cuboidal without goblet cells)

☆ Epithelium: Simple columnar epithelium and Simple cuboidal epithelium with Cilia.

☆ Changes in epithelium is gradual.

In larger bronchioles it's pseudostratified ciliated columnar.
Then gradually changes into simple columnar and simple cuboidal.
Then lose the cilia and change to Clara cells.




Clara cells

- Contain secretory granules: 1- Bactericidal
2- for surfactant

- Contain neuroepithelial bodies

- Receives cholinergic nerve endings → chemoreceptors for changes in O_2 and CO_2 .

- Seen in terminal bronchioles and beginning of respiratory bronchioles.



Terminal Bronchioles



- Closed bronchiole : circular lumen , Rounded
- There are foldings of mucosa because the lamina propria contain smooth muscles and elastic fibers.
- spiral circular smooth muscle.

↳ Contraction of these muscles → Asthma

Sympathetic → Relaxation of the muscle


parasympathetic → Contraction

→ In cases of emergencies we may use Adrenaline.



Respiratory Bronchioles


- Opened bronchiole : opens in an alveolar duct
- lumen is not circular.
- Epithelium : Simple cuboidal ciliated or non-ciliated (clara).
When going distally it become simple squamous epithelium.
- Smooth muscles : knobs and tags of smooth muscles located at the opening of the alveolar duct.
- less folding in mucosa , because of less smooth muscles.
- Between alveoli there are elastic fibers and reticular fibers for inflation and deflation and support for Alveolar duct and Alveoli so it doesn't rupture.
- Alveolar ducts open into Atria that communicate with the alveolar sac.



Alveoli

★ About 300 million Alveoli in lungs.


- Sac-like evagination.
- Diameter: 200 μ m.
- Spongy and soft filled with air and surrounded by elastic fibers



- Between alveoli there are **Interalveolar Septum**
- The Alveolus membrane is made of two types of cells:
 type 1: 97% and type 2: 3% → type 2 is known as **Septal cells**.
- While in the Interalveolar septum, **Septal cells**.
 type 1: 8% and type 2: 16% → mainly in the corners between the alveoli
- Endothelial cells present in the septum and the alveolar membrane. → **Leukocytes + macrophages + fibroblasts**
- Interstitium: CT + capillaries.
- Macrophages: Dust cells
- Between adjacent alveoli there are openings called **Alveolar pores** (for spreading oxygen between the alveoli for equalize the pressure.
 → 10-15 μ m in diameter.

Components of Intra alveolar Septum; **★ Make sure that you can differentiate between the Alveolar membrane and intraalveolar septum.**

- 1- Cells in interstitium
- 2- Endothelial cells (capillaries) 30%
- 3- Fibroblast and Mast cells 36%
- 4- Macrophages 10%
- 5- Type I cells 8%
- 6- Type II cells 16%
- 7- leukocytes



Copyright ©2006 by The McGraw-Hill Companies, Inc.

Blood - Air Barrier


* Also known as respiratory membrane.

* Between the capillary and type I Alveolar cells for gas exchange.
layers:

- 1- Surfactant
- 2- Cytoplasm of alveolar cells
- 3- Fused basal lamina of the closely posed Alveolar and Endothelial cells.
- 4- Cytoplasm of endothelial cells



Total thickness: 0.1 - 1.5 μm .

- Internal Exchange Surface area is approximately 140 m^2 .




Capillary Endothelial Cells

- Extremely thin
- Confused with type I alveolar epithelial cells (both are simple squamous)
 - ↳ under microscope we differentiate endothelial cells by the presence of RBCs.
- Non-fenestrated.
- Clustering of the nuclei and other organelles.
 - And the remanant of cytoplasm is empty to contain O_2 .
- Numerous pinocytic vesicles.





Type I cells.

- Simple squamous epithelium
- 25 nm.
- Cluster of nucleus and organelles and leaving large areas of cytoplasm virtually free.
- Contain pinocytic vesicles.
- There are Dismosomes between type I and type II for occluding junction.



Type II cells

- Cuboidal cells.
- larger than type I
- Secrete surfactant.
- High mitotic activity and able to replace both: type I and II
- Contain lamellar bodies → contain the surfactant.
→ foamy appearance for the cytoplasm.



Pulmonary Surfactant

- 4 types of proteins A, B, C and D.
- Decrease surface tension + Bactericidal effect
- Form the bronchoalveolar fluid.

Alveolar-Lining Regeneration

- Inhalation of NO_2 destroys type I and II.
- Type II turnover rate is high and is able to give both type I and II.

Lung Macrophages

- Dust cells.
- Seen in lungs and may be seen on the surface of placenta.
- Appear as black dots
- Originate from monocytes.
- May go through bronchioles to bronchi then to pharynx. Then undergo expectoration with saliva.
- Huge number exceeds type I cells.

Alveolar (lung) macrophages are more abundant in number than type I alveolar cells.


Type I alveolar cells:

Cover ~95% of the alveolar surface area, but they are few in number (large, flat cells).

Alveolar macrophages:

Are numerous, freely present in the alveolar lumen and septa, and can increase further during inflammation.

So, although type I cells dominate surface area, macrophages dominate cell number.



Pulmonary lymphatic vessels of lung

Superficial and deep lymphatic vessels that all go to the hilum (mediastinal lymph nodes) → lymphatic duct in left side and Right lymphatic duct in right side.
→ Beginning of brachiocephalic vein on left and Right.


Innervation of lung

Lung + visceral pleura



- Sympathetic → bronchodilatation
- parasympathetic → bronchoconstriction

Pleura


- Visceral and parietal
- Contain elastic and reticular fibers
- Innervation like lung sympathetic and parasympathetic.
- Epithelium: Mesothelium (simple squamous epithelium)






Trachea




Extra -
Pulmonary
bronchi



Intrapulmonary
bronchi



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.