


# The Nervous System


- A network of billions of nerve cells linked together in a highly organized fashion to form the rapid control center of the body.



# Basic Functions of the Nervous System

## 1. Sensation

- Monitors changes/events occurring in and outside the body. Such changes are known as *stimuli* and the cells that monitor them are *receptors*.




## 2. Integration

- The parallel processing and interpretation of sensory information to determine the appropriate response

## 3. Reaction

- Motor output.
  - The activation of muscles or glands (typically via the release of neurotransmitters (NTs))




# Nervous Tissue

- Highly cellular
- 2 cell types
  1. Neurons
    - Functional, signal conducting cells
    - Do not divide
    - Long lived
    - High metabolic activity
    - Electrically excitable
  2. Neuroglia
    - Support, nourish, and protect neurons
    - Divide
    - Smaller cells but they greatly outnumber neurons by about 5 to 50
    - 6 types of supporting cells: (4 are found in the CNS, and 2 are found in the PNS.)

- **White matter:** aggregations of myelinated and unmyelinated axons of many neurons
- **Gray matter:** contains neuronal cell bodies, dendrites, unmyelinated axons, axon terminals, and neuroglia


  

- **Nerves:** Bundles of processes in the PNS
- **Tracts:** Bundles of processes in the CNS (No Connective tissue)
- **Ganglion:** cluster of nerve cell bodies in PNS
- **Nucleus:** cluster of nerve cell bodies in CNS (surrounded by white matter)
  - If not surrounded (Cortex)






- A bundle of processes in the PNS is a **nerve**.
- Within a nerve, each axon is surrounded by an **endoneurium**
- Groups of fibers are bound together into bundles (fascicles) by a **perineurium**
- All the fascicles of a nerve are enclosed by a **epineurium**



# Organization of the Nervous System

- Anatomical divisions:
  1. Central Nervous System →
    - The brain + the spinal cord
      - The center of integration and control
  2. Peripheral Nervous System
    - The nervous system outside of the brain and spinal cord
    - Consists of:
      - 31 Spinal nerves
        - Carry info to and from the spinal cord
      - 12 Cranial nerves
        - Carry info to and from the brain





# Brain


- **Forebrain:**  
(Prosencephalon)

- Cerebrum:  
(Telencephalon)
- Diencephalon
  - Thalamus
  - Hypothalamus
  - Epithalamus
  - Subthalamus

- **Midbrain:**  
(Mesencephalon)


- **Hindbrain:**  
(Rhombencephalon)

- Pons
- Medulla oblongata
- Cerebellum




# External anatomy of Spinal Cord

- Runs through the vertebral canal
- Extends from foramen magnum to second lumbar vertebra
- Regions
  - Cervical (8)
  - Thoracic (12)
  - Lumbar (5)
  - Sacral (5)
  - Coccygeal (1)
- Gives rise to (31) pairs of spinal nerves
  - All are *mixed* nerves
- Not uniform in diameter
  - Cervical enlargement: supplies upper limbs
  - Lumbar enlargement: supplies lower limbs




# External anatomy of Spinal Cord

- Flattened slightly anteriorly and posteriorly
- length of the adult spinal cord ranges from 42 to 45 cm
- Conus medullaris- tapered inferior end (conical structure)
  - Ends between L1 and L2
- Cauda equina - origin of spinal nerves extending inferiorly from conus medullaris.




# Meninges

- Connective tissue membranes
  - **Dura mater:**
    - Outermost layer; continuous with epineurium of the spinal nerves
    - Dense irregular connective tissue
    - from the level of the foramen magnum to S2
    - Closed caudal end is anchored to the coccyx by the **filum terminale externum**
  - **Arachnoid mater:**
    - Thin web arrangement of delicate collagen and some elastic fibers.
    - Adheres to the inner surface of the dura mater




# Meninges

- Connective tissue membranes
- Pia mater:
  - Bound tightly to surface
  - Thin transparent connective tissue layer that adheres to the surface of the spinal cord and brain
  - Forms the filum terminale
    - anchors spinal cord to coccyx
  - Forms the denticulate ligaments that attach the spinal cord to the arachnoid mater and inner surface of the dura mater



# Spaces


- **Epidural:** space between the dura mater and the wall of the vertebral canal.
  - Anesthetics injected here
  - Fat-fill
- **Subdural space:** serous fluid
- **Subarachnoid:** between pia and arachnoid
  - Filled with CSF
  - Lumbar puncture
  - supracristal line
  - L3-L4



# Spinal cord segment


- The segments of the spinal cord are not in line with the corresponded vertebrae and the difference increases as we go downward.
- The roots increase in length as you go downward.
- Every spinal nerve emerges from the spinal column through the intervertebral foramen under its corresponding vertebra
- first 7 cervical nerves pass above their corresponding vertebrae

Spinous process	spinal cord segment
C7	C8
T3	T5
T9	T12
T10	L1-2
T11	L3-4
T12	L5
L1	S1-end




# Herniated Disc/ ruptured disc/ slipped disc



protrusion (leakage) of the gelatinous nucleus pulposus through the anulus fibrosus of IV disc



**Posterolateral direction:**  
**Thinner annulus fibrosus**

**95% in L4/L5 or L5/S1**






# Common lumbar disc problems

Disc	Root	Percentage	Motor weakness	Sensory changes	Reflex affected
L3-L4	L4	3-10%	Knee extension (Quadriceps femoris)	Anteriomedial leg (saphenous)	Knee jerk
L4-L5	L5	40-45%	Big toe dorsiflexion (EHL) and TA	Big toe , anteriolateral leg (Common P)	Hamstring jerk
L5-S1	S1	45-50%	Foot planter flexion (Gastrocnemius)	Lateral border of foot (sural)	Ankle jerk

## Important myotomes of lower limb

□ **Test L5:** by asking the patient to stand on his heels


□ **Test S1:** by asking the patient to stand on his tiptoes




(C) Anterior View




(D) Medial View



(B) Lateral View



**CERVICAL SPINE**



Both discs affect same nerve root in cervical spine, different than lumbar spine

Lumbar Spine Pedicle/nerve Root  
MATCH

C8 and above Pedicle / Nerve Root  
MISMATCH


"Extra C8 Nerve Root (without C8 pedicle)  
allows transition from MISMATCH to MATCH

T1 and below Pedicle / Nerve Root  
MATCH




# Major symptoms of disc herniation

- **Low back pain:** radiating to the gluteal region, the back of the thigh and back of the leg
- spinal nerve gives a meningeal branch bring sensation from the dura matter
- Dura matter is sensitive to stretch
- Pain is diffused due to overlapping dermatomes
- **Straight Leg Raise Test (SLR)**




- ❑MRI is commonly used to aid in making the diagnosis of a herniated disc





# Cross Section of Spinal Cord

- Anterior median fissure:  
wide groove on the Anterior aspect
- posterior median sulcus:  
Narrow groove on the posterior aspect
- Gray matter: neuron cell bodies, dendrites, axons
  - Divided into *horns*
  - **Posterior (dorsal) horn**  
(cell body of sensory N)
  - **Anterior (ventral) horn**  
(cell body of motor N to skeletal M)
  - **Lateral horn**  
(cell body of motor N to cardiac M, smooth M, glands)




# Cross Section of Spinal Cord






# Posterior White Column-Medial Lemniscal Pathway

- Modality: Discriminative Touch Sensation (include Vibration) and Conscious Proprioception
- Receptor: Most receptors except free nerve endings
- 1st Neuron: Dorsal Root Ganglion
- 2nd Neuron: Dorsal Column Nuclei (Nucleus Gracilis and Cuneatus)
  - Internal Arcuate Fiber - Lemniscal Decussation
  - Medial Lemniscus
- 3rd Neuron: Thalamus (VPL) Internal Capsule ----- Corona Radiata
- Termination: Primary Somesthetic Area (S I)



# Posterior White Column-Medial Lemniscal Pathway



Discriminative touch, vibratory sense, and conscious muscle-joint sense

• **Posterior Column tract consists of:**


• **Fasciculus gracilis**

• Transmits information coming from areas inferior to T6


• **Fasciculus cuneatus**

• Transmits information coming from areas superior to T6


Figure 15.1 Anatomical Principles for the Organization of the Sensory Tracts and Lower-Motor Neurons in the Spinal Cord




## Discriminative Touch Spinal Pathway




## Posterior Columns





## Primary Somatosensory (SI) Cortex


- Axons from third-order thalamic neurons terminate in the **primary somatosensory (SI) cortex**
- subdivided into four distinct areas; from anterior to posterior, these are **Brodmann areas 3a, 3b, 1, and 2**



- **Area 3a:** muscle spindle afferents (mainly)
- **Area 2:** Golgi tendon organs, and joint afferents (mainly).
- **Areas 3b and 1:** They receive cutaneous afferents from receptors such as Meissner corpuscles and Merkel cells). also receive input from cutaneous receptors that transmit pain and temperature

# **lateral spinothalamic tract**

- Modality: pain and temperature
- Receptors: free nerve endings
- 1<sup>st</sup> Neuron: Dorsal root ganglia
- 2<sup>nd</sup> Neuron: the posterior gray column (substantia gelatinosa)  
The axons of 2<sup>nd</sup> order neurons cross obliquely to the opposite side in the anterior gray and white commissures , ascending in the contralateral white column as the lateral spinothalamic tract
- 3<sup>rd</sup> Neuron: Thalamus (VPL)  
Internal Capsule ----- Corona Radiata
- Termination: Primary Somesthetic Area (S I) and Widespread Cortical Region



# Rexed laminae

- **Lamina 1** relay information related to pain and temperature
- **Lamina 2:** relay information related to pain and temperature **(pain modulation)**
- **Lamina 3 and 4:** nucleus proprius; these laminae have many interneurons






Fig. 5.2. Subdivisions of the grey matter of the spinal cord. The left half of the figure shows the cell groups usually described. The right half shows the newer concept of laminae.

- **Lamina 5:** relay information related to pain and temperature
- **Lamina 6:** presents only at the cervical and lumbar enlargements and receives proprioception
- **Lamina 7:** **Intermedio-lateral** nucleus, contains preganglionic fibers of sympathetic (T1 -L2). **Intermedio-medial nucleus**, all over the spinal cord, receive visceral pain. **Dorsal nucleus of Clark's** presents at (C8 – L2 or T1- L4) , relay center for **unconscious proprioception**

# lateral spinothalamic tract



A





B

Source: Ropper AH, Samuels MA, Klein JP; Adams and Victor's Principles of Neurology, Tenth Edition: [www.accessmedicine.com](http://www.accessmedicine.com)  
Copyright © The McGraw-Hill Companies, Inc. All rights reserved.

- Lamina 1+ 5: the spinothalamic tract ascend which transmit pain, temperature and touch. (A delta fibers)
- Lamina 1+ 2: the spinothalamic tract ascend (C fibers).


## Posterolateral tract of Lissauer

- located between the posterior white column and the lateral white column




# Other Terminations of the Lateral Spinothalamic Tract


- **Reticular formation:** (majority of the slow pain fibers) individual becomes aware of the pain
- **Cingulate gyrus:** interpretation of the emotional aspect of pain
- **Insular gyrus:** concerned with the interpretation of pain stimuli from the internal organs of the body and brings about an autonomic response






# Referred pain mechanism convergence theory

- ❑ Referred pain is presumed to occur because the information from multiple nociceptor afferents converges onto individual spinothalamic tract neurons
- ❑ The brain therefore interprets the information coming from visceral receptors as having arisen from receptors on the body surface, since this is where nociceptive stimuli originate more frequently









# Anterior spinothalamic tract

- Modality: crude touch and pressure
- Receptors: free nerve endings
- 1<sup>st</sup> Neuron: Dorsal root ganglia
- 2<sup>nd</sup> Neuron: the posterior gray column (**nucleus proprius**)  
The axons of 2<sup>nd</sup> order neurons cross obliquely to the opposite side in the anterior gray and white commissures , ascending in the contralateral white column as the Anterior spinothalamic tract
- 3<sup>rd</sup> Neuron: Thalamus (VPL)  
Internal Capsule ----- Corona Radiata
- Termination: Primary Somesthetic Area (S I)



# Spinotectal Tract


- ascend in the anterolateral white column lying close to the lateral spinothalamic tract
- Terminate: superior colliculus
- Provides afferent information for spinovisual reflexes



**In Medulla:** ant spinothalamic tract + spinotectal + lateral spinothalamic = spinal lemniscus

# Posterior spinocerebellar

- muscle and joint sensation
- 1<sup>st</sup> order neuron axons terminate at the base of post gray column (nucleus dorsalis or **Clarks nucleus**)
- the axons of 2<sup>nd</sup> order neurons enter posterolateral part of the lateral white matter **on the same side**
- ascend as the posterior spinocerebellar tract to medulla oblongata
- Terminates in cerebellar cortex (through inferior cerebellar peduncle)



➤ note: axons of lower lumbar and sacral spinal nerves ascend in the posterior white column until they reach L3 or L4 segments where they synapse with nucleus dorsalis

# Rexed laminae

- **Lamina 1** relay information related to pain and temperature
- **Lamina 2:** relay information related to pain and temperature **(pain modulation)**
- **Lamina 3 and 4:** nucleus proprius; these laminae have many interneurons





Fig. 5.2. Subdivisions of the grey matter of the spinal cord. The left half of the figure shows the cell groups usually described. The right half shows the newer concept of laminae.


- **Lamina 5:** relay information related to pain and temperature
- **Lamina 6:** presents only at the cervical and lumbar enlargements and receives proprioception
- **Lamina 7:** **Intermedio-lateral** nucleus, contains preganglionic fibers of sympathetic (T1 -L2). **Intermedio-medial nucleus**, all over the spinal cord, receive visceral pain. **Dorsal nucleus of Clark's** presents at (C8 – L2 or T1- L4) , relay center for **unconscious proprioception**

# Anterior spinocerebellar tract

- muscle and joint sensation
- 1<sup>st</sup> order neuron axons terminate at the base of post gray column (nucleus dorsalis)
- the majority of axons of 2<sup>nd</sup> order neurons cross to opposite side and ascend as anterior spinocerebellar tract in the contralateral white column
  - *the minority of axons ascend as anterior spinocerebellar tract in the lateral white column Of the same side*
- ascend as anterior spinocerebellar tract to medulla oblongata and pons
- Terminates in cerebellar cortex (through superior cerebellar peduncle)
  - *the fibers that crossed over in spinal cord cross back within cerebellum*




## Spinocerebellar Tracts



Proprioceptive input from Golgi tendon organs, muscle spindles, and joint capsules

# Motor tracts

- There are two major descending tracts
- **Pyramidal tracts**  
(Corticospinal ) : Conscious control of skeletal muscles
- **Extrapyramidal:**  
Subconscious regulation of balance, muscle tone, eye, hand, and upper limb position:
  - ❖ **Vestibulospinal tracts**
  - ❖ **Reticulospinal tracts**
  - ❖ **Rubrospinal tracts**
  - ❖ **Tectospinal tracts**



Extrapyramidal tracts arise in the brainstem, but are under the influence of the cerebral cortex

# Rexed laminae

- **Lamina 8:** motor interneurons, Commissural nucleus
- **Lamina 9:** ventral horn, LMN, divided into nuclei:
  - **Ventromedial:** all segments (extensors of vertebral column)
  - **Dorsomedial:** (T1-L2) intercostals and abdominal muscles
  - **Ventrolateral:** C5-C8 (arm) L2-S2 (thigh)
  - **Dorsolateral:** C5-C8 (Forearm), L3-S3 (Leg)
  - **Reterodorsolateral:** C8-T1 (Hand), S1-S2 (foot)
  - **Central:** Phrenic nerve (C3-C5)
- **Lamina X:** Surrounds the central canal – the grey commissure





Fig. 5.2. Subdivisions of the grey matter of the spinal cord. The left half of the figure shows the cell groups usually described. The right half shows the newer concept of laminae.

Figure 15.1 Anatomical Principles for the Organization of the Sensory Tracts and Lower-Motor Neurons in the Spinal Cord

MEDIAL

LATERAL



© 2012 Pearson Education, Inc.

- Motor neurons of anterior horn
- **Medial group:** (All segments)
- **Lateral group:** only enlargements


# Brain stem

- Stalk like in shape
- Connects spinal cord and forebrain


Parts:

1. Medulla oblongata
2. Pons
3. Midbrain

## Brain Stem



Anterior view



# Motor tracts


□ Both pyramidal tracts and extrapyramidal both starts from cortex:

- Area 4
- Area 6
- Area 312

❖ Pyramidal: mainly from area 4


❖ Extrapyramidal: mainly from area 6

- area 6
- **Premotor area:** uses external cues
- **Suplementary motor area:** uses internal cues





# Lateral corticospinal tract

- The upper motor neurons of these tracts originate in the precentral gyrus of the cerebral cortex
- In midbrain:middle three-fifths of the **basis pedunculi of the midbrain**
- In medulla oblongata: pyramids
- Most of the fibers (85 percent) cross over (decussate) to the opposite side in the pyramidal decussation, where they continue to descend in the **lateral funiculus** of the spinal cord as the **lateral corticospinal tract (LCST)**.



- The tract descends all the way of spinal cord with fibers continually leaving it in order to synapse on interneurons in the anterior gray horn. ( Some even synapse directly on alpha and gamma motor neurons)
- *Those corticospinal fibers which do not decussate in the medulla continue descending on the same (ipsilateral) side of the cord and become the anterior corticospinal tract (ACST).*







# corticospinal tract

for fine skilled movements

- Lateral corticospinal tract descends the full length of the spinal cord
- LCST fibers synapse with alpha and gamma nuclei of the
  - Cervical region (55%) (great effect on the upper limb)
  - Thoracic 20%
  - Lumbar and Sacral 25%
- The lateral corticospinal tract synapses **mainly by interneurons** In laminae IV, V, VI, VII, VIII
- **Exception:** 3% originate from the fifth layer of area 4 (giant cells of betz) synapse directly. (Accurate movements)





## The anterior corticospinal tract

acts on the proximal muscles of upper limb (shoulder muscle) of the ipsilateral and contralateral sides


*Fibers leave the tract at various levels to cross over in the anterior white commissure to synapse on interneurons in the anterior gray horn.*

## The Corticoneuclear Tract (fibers)

- This tract is composed of fibers originating in the precentral gyrus of the lower quarter of the motor cortex.
- The descending fibers terminate in the motor nuclei of cranial nerves III and IV in the midbrain; V, VI, and VII in the pons; and IX, X, XI, and XII in the medulla.
- The corticobulbar fibers from one side of the brain project to the motor nuclei on both sides of the brainstem (bilateral input)

The corticoneuclear input is bilateral **Except :**

- 1- Part of 7<sup>th</sup> ( which supplies LOWER facial muscles)
- 2- Part of 12<sup>th</sup> (which supplies genioglossus muscle)



## **The Subconscious Motor Tracts**

- Consists of four tracts involved in monitoring the subconscious motor control
- **Vestibulospinal tracts**
- **Tectospinal tracts**
- **Reticulospinal tracts**
- **Rubrospinal tracts**


Extrapyramidal tracts arise in the brainstem, but are under the influence of the cerebral cortex

These motor pathways are complex and multisynaptic, and regulate:

- Axial muscles that maintain balance and posture
- Muscles controlling coarse movements of the proximal portions of limbs
- Head, neck, and eye movement

# Rubrospinal tract


- **Red nucleus**
  - In the midbrain at the level of superior colliculus
  - Receives afferent fibers from cerebral cortex and the cerebellum
- **Crossed** (at the level of the nucleus)
- Lateral white column
- **Function:**  
facilitate the activity of flexors and inhibit the activity of extensors



# Rubrospinal tract


- rubrospinal tract is very close to the lateral corticospinal tract in the spinal cord. They form the **lateral motor system**
- synapses with alpha and gamma through interneurons
- Excitatory to flexors and inhibitory to extensors
- supply the distal flexors muscles mainly with little effect on the proximal muscles

( facilitate the activity of flexor muscles )




## Pontine reticulospinal tract

- From pons:
- axons of RF neurons descend **uncrossed** into the spinal cord
- Anterior white column
- medial reticulospinal tract (MRST)
- **tonically active**
- normally under **inhibition from cortex**
- **Function:**
- activate the axial and proximal limb extensors



# Medullary reticulospinal tracts


- From medulla
- axons of RF neurons descend **crossed and uncrossed** into the spinal cord
- Lateral white column
- Lateral reticulospinal tract (LRST)
- NOT tonically active
- normally under **stimulation**
- **Function:**  
Inhibit the axial and proximal limb extensors



# Vestibulospinal Tract

- **Vestibular nuclei**
  - in the pons and medulla beneath the floor of 4<sup>th</sup> ventricle
  - Receives afferent fibers from the inner ear through the vestibular nerve and from the cerebellum
- **Uncrossed**
- Anterior white column
- **Function:**

facilitate the activity of extensor muscles and inhibit the activity of flexor muscles association with the maintenance of balance



# Vestibulospinal tract

- nerve cells in vestibular nucleus (in the pons and medulla oblongata)
  - received afferents from inner ear and cerebellum
- axons descend uncrossed
  - through medulla and through the length of spinal cord
- synapse with neuron in the anterior gray column of the spinal cord

( balance by facilitate the activity of the extensor muscles )






Fig. 5.19 Vestibulospinal tracts.



# Reticulospinal tracts


- Has also descending autonomic fibers providing a pathway by which the hypothalamus can control the sympathetic and sacral parasympathetic outflow.
- Most of these fibers are derived from *the lateral reticulospinal tract*



# Tectospinal tract

- nerve cells in superior colliculus of the midbrain
- **Crossed**
- The tract descends in the anterior white column close to Anterior median fissure
- Majority of fibers terminate in the anterior gray column of upper cervical segments of spinal cord

( responsible for reflex movement of head & neck in response to visual stimuli )



# The motor pathways are classified into

- **Medial Motor system:** axial & proximal muscles. Medial Motor system include:
  - Anterior corticospinal tract.
  - Extrapyramidal pathway in general
- **Lateral Motor system:** distal muscles mainly, lateral Motor system include
  - lateral corticospinal tract
  - Rubrospinal tract distal muscles mainly (and proximal).