

---

# **Cerebral Cortex; Intellectual Functions of the Brain; Learning and Memory-**

---

Faisal I. Mohammed, MD, PhD

---

# Objectives


- Describe intellectual functions of the cerebral cortical areas
- Explain memory and learning
- Outline the dominant and non-dominant hemispheres
- Delineate language areas of speech

---

# Physiologic Anatomy of Cerebral Cortex

- Each area of the cortex is connected to a specific part of the thalamus.
  - When thalamic connection is lost cortical function stops.
  - All sensory pathways pass through the thalamus with the exception of some olfactory signals.
-

Frontal plane



DURA MATER:

Periosteal layer

Meningeal layer

Subarachnoid space

Arachnoid villus

FALX CEREBRI

Superior sagittal sinus

Skin

Parietal bone cranium

CRANIAL MENINGES:

Dura mater


Arachnoid


Pia mater

Cerebral cortex


Frontal section through skull showing the cranial meninges

14.02






(a) Frontal section of primary somatosensory area in right cerebral hemisphere



(b) Frontal section of primary motor area in right cerebral hemisphere


# Functional organization of the primary Motor Cortex




(b) Frontal section of primary motor area in right cerebral hemisphere

- ❖ Located in the precentral gyrus of the frontal lobe.
- ❖ More cortical area is devoted to those muscles involved in skilled, complex or delicate movements, that have more motor units i.e the cortical representation is proportional to the No of motor units

# Physiological Anatomy of the Cerebral Cortex





# Functional Organization of the Cerebral Cortex: Sensory Areas



Lateral view of right cerebral hemisphere

# Functional Areas of the Cerebral Cortex





---

# Dominant and Non-Dominant Hemisphere

- Wernicke's area more developed in one hemisphere, responsible for verbal symbolism and related intelligence. *Dominance is related to Language*
  - 95% of population has a left dominant hemisphere.
  - Wernicke's area can be as much as 50% larger in the dominant hemisphere.
-

---

# Dominant and Non-Dominant Hemisphere (Cont'd)

- Damage to dominant Wernicke's area leads to dementia.
  - Non-dominant side related to other forms of sensory intelligence (music, sensory feelings).
-

# Hemispheric Lateralization

**TABLE 14.3**

## Functional Differences Between the Two Cerebral Hemispheres

### LEFT HEMISPHERE FUNCTIONS

Receives somatic sensory signals from and controls muscles on right side of body.

Reasoning.

Numerical and scientific skills.

Ability to use and understand sign language.

Spoken and written language.

### RIGHT HEMISPHERE FUNCTIONS

Receives somatic sensory signals from and controls muscles on left side of body.

Musical and artistic awareness.

Space and pattern perception.

Recognition of faces and emotional content of facial expressions.

Generating emotional content of language.

Generating mental images to compare spatial relationships.

Identifying and discriminating among odors.

Table 14.03 Tortora - PAP 12/e  
Copyright © John Wiley and Sons, Inc. All rights reserved.

# Language Areas

- Located in a large area surrounding the left (or language-dominant) lateral sulcus
- Major parts and functions:
  - Wernicke's area – involved in sounding out unfamiliar words – sensory aspect of speech- damage → sensory aphasia (Receptive aphasia)
  - Broca's area – speech preparation and production- motor aspect of speech – damage → motor aphasia (expressive aphasia)
  - Both Wernicke's and Broca's area damage → global aphasia
  - Lateral prefrontal cortex – language comprehension and word analysis
  - Lateral and ventral temporal lobe – coordinate auditory and visual aspects of language

# Brain Organization and Handedness

- Close to 90% of people are right-handed and close to 10% are left-handed and a small number are ambidextrous (use both hands)
- 95% of right-handers process speech primarily in the left hemisphere
  - left-handers: around 65% in left hemisphere, 15-20% in right hemisphere, 15-20% in both
- More than 90% of people are born with the left hemisphere area that controls the movement of the right hand is bigger
  - They tend to use the right hand, this area grows and become dominant

---

# Brain Organization and Handedness

## ...cont

- Left handed people have their right cerebral hemisphere area that controls the movement of the left hand bigger
  - If they use the left hand then this area grow and become dominant
  - They still can convert and the younger the more easier if they start to use their right hand instead and then they become right handed
  - Same applies for using the legs

---

# Intellectual Functions of the Prefrontal Association Area

- responsible for calling forth stored information and using it to obtain a goal
  - responsible for concerted thinking in a logical sequence
    - damage causes an inability to keep track of simultaneous bits of information, easily distracted
-

---

# Intellectual Functions of the Prefrontal Association Area (Cont'd)

- elaboration of thought
    - prognosticate, plan, consider consequences of motor actions before they are performed
    - correlate widely divergent information, control one's activities
    - Personality trait and behavior that confines to values and manners of the culture
-

# Pathways for Auditory Communication

5. activation of motor programs in Broca's area for control of word formation


6. transmission of signals to motor cortex to control speech muscles

4. transmission via the arcuate fasciculus to Broca's area


3. formation of the word that expresses a particular thought

2. interpretation of the word and the thought that the word expresses in Wernicke's area

1. primary auditory area recognition of the sound as a word



## Speaking a heard word



# Pathways for Visual Communication


5. transmission of signals to motor cortex to control speech muscles

4. then to Broca's area for motor formation of the word


3. visual input reaches full level of interpretation in Wernicke's area

2. processing of the visual information in the parietal-temporal-occipital association cortex, the angular gyrus region


1. receive the visual input in primary visual area



## Speaking a written word



# Sensory Aspects of Communication



## Wernicke's aphasia

Destruction of the visual and auditory association areas results in an inability to understand the written or spoken word.

---

# Motor Aspects of Communication

- Speech involves two things
    - formation in the mind of thoughts to be expressed and the choice of words
    - motor control of vocalization and the act of vocalization
  - Formation of word, thought and choice of words is function of Wernicke's area.
  - Broca's area controls the motor coordination required for speech.
-

---

# Function of the Corpus Callosum

- connects the two hemispheres and allows transfer of information
  - interruption of these fibers can lead to bizarre types of anomalies
    - dominant hemisphere understands spoken word
    - non dominant hemisphere understands written word and can elicit motor response without dominant side knowing why response was performed
-

---

# Thoughts and Memory

- Neural mechanism for thought is not known.
  - Most likely a specific pattern of simultaneous neural activity in many brain areas.
  - Destruction of cerebral cortex does not prevent one from thinking.
    - However, depth of thought and level of awareness may be less.
-

---

# Memory and learning

- Learning is acquiring new sensory information or motor skills
  - Change in the capability of synaptic transmission from neuron to neuron as a result of prior stimulation.
  - Memory trace is a specific pattern or pathway of signal transmission.
  - Once established they can be activated by the thinking mind to reproduce the pattern and thus the memory.
-

---

# 3 Types of Memory

- immediate memory
    - lasts for seconds or minutes (remembering 10 digit phone number)
  - short-term memory
    - lasts for days to weeks
  - long-term memory
    - lasts for years or for a lifetime
-

---

# Mechanism of Memory

- Immediate memory may result from synaptic potentiation through the accumulation of calcium in the presynaptic membrane.
    - would promote neurotransmitter release
  - Short-term memory may result from a temporary physical or chemical change in the pre- or postsynaptic membrane.
-

---

# Cellular Basis for Memory

- repetitive stimulation causes a progressive decline in sensitivity called *habituation*
  - results from progressive decline in the number of active calcium channels
  - less calcium entry less transmitter released
  - stimulation of facilitator terminal prevents habituation
-

# Molecular Basis for Memory

Transmitter activates G protein which in turn activates adenylate cyclase resulting in an increase in cAMP.

cAMP activates a protein kinase that phosphorylates a component of the  $K^+$  channel blocking its activity.

This prolongs the action potential which increases transmitter release.




Figure 57-9; Guyton & Hall

---


# Long-Term Memory

- results from a structural change in the synapse
  - increase in the area for vesicular release  
therefore, more transmitter is released
  - during periods of inactivity the area decreases  
in size
  - enlargement of the release site area results  
from synthesis of release site proteins
-

---

# Consolidation of Memory

- converting immediate into short and long-term memory
  - results from chemical, physical and anatomical changes in the synapse
  - requires time
  - interruption of the process by electrical shock or by anesthesia will prevent memory development
  - rehearsal enhances consolidation
-




---

# Brain Centers and Memory

- Thalamic structures are important for recalling memories.
  - Damage to thalamus causes **retrograde amnesia** or the inability to recall stored experiences.
  - Thalamus scans the cortex for the area and the circuit for the stored memory.
-

# Comparison of Long-Term and Short-Term Memory

CHARACTERISTIC	SHORT-TERM MEMORY	LONG-TERM MEMORY
<b>Time of Storage after Acquisition of New Information</b>	Immediate	Later; must be transferred from short-term to long-term memory through consolidation; enhanced by practice or recycling of information through short-term mode
<b>Duration</b>	Lasts for seconds to hours	Retained for days to years
<b>Capacity of Storage</b>	Limited	Very large
<b>Retrieval Time (remembering)</b>	Rapid retrieval	Slower retrieval, except for thoroughly ingrained memories, which are rapidly retrieved
<b>Inability to Retrieve (forgetting)</b>	Permanently forgotten; memory fades quickly unless consolidated into long-term memory	Usually only transiently unable to access; relatively stable memory trace
<b>Mechanism of Storage</b>	Involves transient changes in functions of pre-existing synapses, such as altering amount of neurotransmitter released	Involves relatively permanent functional or structural changes between existing neurons, such as formation of new synapses; synthesis of new proteins plays key role



# Thank You

