



Preoperative Assessment

Dr. Ahmad Arraqap
Anesthesia dept.
Faculty of Medicine
University of Jordan

Learning Objectives

- After attending this lecture, student is expected to be to:

1. Aims of preoperative assessment.
2. How preoperative assessment is conducted.
3. ASA risk score.
4. Fasting guidelines.
5. Preoperative preparation of patients before anesthesia.

Aims of preoperative assessment

- Opportunity to identify **co-morbidities** that may lead to patient **complications** during the **peri**-operative period.
- Optimize any co-morbidities.
- **Venue**: Preoperative clinic or Wards (anaesthesia clinic) for elective cases or Emergency Department/ward for emergent surgeries.
- Establish a rapport.



Conduct of Assessment

- **History, Physical** examination, and **Investigations** +/- further consultations.
- Make sense of collected data to formulate **anaesthetic plan**.



PHYSICAL ASSESSMENT

COLLECT OBJECTIVE DATA

* USING SENSES



RASH



WHEEZES

VALIDATE SUBJECTIVE INFORMATION

* HEALTH HISTORY



DEVELOP PLAN of CARE



EVALUATE
EFFECTIVENESS of
INTERVENTION



History

- **Profile:**
 - Name/ Age/ Gender/ Weight/ Height
 - Type of surgery
 - Smoking history
 - Fasting hours
- **Review of Systems (focused):**

Cardiovascular

IHD (CP /Angina/ stent)
CHF (PNDs/ orthopnea)
Exercise intolerance
Palpitations

Respiratory

Asthma
COPD
OSA
Recent URTI/LRTI
Cough/ sputum
Smoking

Neurologic

-Epilepsy
-CVA/TIA
-Denervation disease

GIT

GERD
PUD
Hiatus Hernia
Intestinal obstruction.

Renal

CRF
ARF
On dialysis

Blood disorders
Antiplatelet
Anticoagulation


History

- Past medical history.
- Medications including allergies.
- Surgical history including previous anaesthesia.

- **PREVIOUS ANAESTHESIA:**

- Very important part of previous anaesthesia history is airway related history.
 - Previous difficult airway.
 - Previous airway surgeries/ burns.
 - Snoring/ obstructed breathing.
- Always check previous records/ old file.

- Previous anesthesia
- Type of anesthesia
- Complications: difficult airway management/delayed emergence / PONV)-
- Family hx.



Physical Examination

- **General** appearance: Obesity, malnutrition, pregnancy head and neck ..etc
- **Vital** signs: BP, HR, RR.

- **Cardiac** exam

Rate and rhythm
Heart sound

- **Respiratory** exam

Sign of respiratory distress
Respiratory rate
Auscultate Lung

- **Neuro** exam

Mental status
Gross motor/gross sensory

Airway examination

- Purpose: To anticipate any possible difficulty in ventilation and intubation.
- Importance: airway and respiratory events are the most common events during anaesthesia. (most common: sore throat and dental damage).

Method of Assessment (L.E.M.O.N)

L

Look externally

face / mouth opening/ teeth / tongue

E

Evaluate the three distances

interincisal / thyromental / sternomental distance

M

Mallampati score (3 or 4)

O

Obstruction (presence of any obstruction like:

peri-tonsillar abscess , thyroid mass , VC nodule)

N

Neck mobility

INSPECTION

1. Does it **look** difficult?

- Obesity
- Deformities, masses, scars or burns.
- Large breasts in females.
- Neck deformities or large neck fat pad
- Position of the mandible: excessive protrusion or recession.
- Nasal deformity, deviation, patency of nostrils.
- Mouth asymmetry, deviation, high arched palate, large tonsils, abscess.
- Dentition: protrusion, missing/loose, hygiene, crowns and caps.
- BEARD



INSPECTION

2. *Mouth opening.*

- At least 3 fingers of patient's own.

3. *Mobility of the lower jaw and neck.*

- Ability of protruding the lower jaw in front of the upper one.
- Neck extension and flexion.



Three distances

Thyro-mental distance

- ✓ It describes the distance between the mentum & thyroid notch
- ✓ It helps in determining how readily the laryngeal axis will fall in line with the pharyngeal axis
- ✓ It is normally $> 6\text{cm}$ in adults .

Sterno- mental distance

- ✓ It describes the distance between the mentum & suprasternal notch
- ✓ If this distance less than 12 cm it predicts difficult intubation


Inter-incisor distance

- ✓ It describes the distance between the upper and lower incisors
- ✓ It is normally 4.5 cm

Mallampati score

- Patency of mouth and throat cavities (tongue/mouth).

class	Structures identified when pt seated
-------	--------------------------------------



Mallampati One

1	Tonsillar pillars, Uvula , soft & hard palate
---	--





THE UNIVERSITY OF
JORDAN

Mallampati two

2	Uvula ,soft & hard palate



Mallampati three

3

Base of uvula ,soft & hard palate





Mallampati four

4

Only hard palate is can be seen



PALPATION

- Submandibular and submental area for masses.
- Tracheal centralization.



Preoperative investigations

- **Routine** (test **ordered in the absence** of a specific clinical indication) testing does **not** add much to the preoperative assessment.
- Guided by history, physical examination, and nature of surgery.
- Haemoglobin/haematocrit.
- Urine Analysis.
- Chest radiograph (CXR).
- Electrocardiograph (ECG).
- Pulmonary Function Tests (PFT).





- CBC

Advanced age/ Anemic pt/ Bleeding /chronic disease (kidney liver heart)

- KFT

Diabetics/ HTN/ chronic disease / on medications like diuretics , digoxin ,ACEI

- Sugar

Diabetics / HTN/ chronic disease / on steroid

- LFT

Liver disease / malnourished pt

- Coagulation

Bleeding disorder/ Kidney disease/ Liver disease/ pt on anticoagulants

Investigations

CXR

- ✓ Indicated in
 - patients with respiratory or cardiac disease
 - smokers
 - patients with recent LRTI

ECG

- ✓ Indicated in
 - patients with respiratory or cardiac disease
 - Advanced Age (M: 55y F: 65y)
 - Any patient with CAD risk factors : (HTN, DM, hyperlipidemia , exercise intolerance)

Investigations

Pulmonary Function test:

- Identifying patients at respiratory risk, evaluating the risk, and finding modified factors to decrease risk

Indicated in:

- Obstructive lung disorders
- Restrictive lung disorders
- Neuromuscular disorders

Includes mainly

- ☐ Spirometry
- ☐ ABGs



ASA risk score

Category	Health status	Examples
ASA 1	A normal healthy patient	Nonsmoker, BMI <30
ASA II	A patient with mild systemic disease	No functional limitations and a well-controlled disease (e.g., treated hypertension, obesity with BMI under 35, frequent social drinker, or cigarette smoker)
ASA III	A patient with a severe systemic disease that is not life-threatening	Some functional limitation due to disease (e.g., poorly treated hypertension or diabetes, morbid obesity, chronic renal failure, a bronchospastic disease with intermittent exacerbation, stable angina, implanted pacemaker)
ASA IV	A patient with a severe systemic disease that is a constant threat to life	(e.g., unstable angina, poorly controlled COPD, symptomatic CHF, recent (less than three months ago) myocardial infarction or stroke)
ASA V	A moribund patient who is not expected to survive without the operation	(e.g., ruptured abdominal aortic aneurysm, massive trauma, and extensive intracranial hemorrhage with mass effect)
ASA IV	A brain-dead patient whose organs are being removed with the intention of transplanting them into another patient	

EMERGENCY ?!

ASA: “when the delay in treatment of the patient would lead to a significant increase in the threat to life or body part.”

E

ASA 1 - E

ASA 2 - E

ASA 3 - E

ASA 4 - E

ASA 5 - E

ASA 6 - E

Plan

- Decisions about:
 - Further consultations.
 - Further investigations.
 - Patient's medications.
 - Preparation of blood and blood products.
 - Type of admission (if seen in clinic)
 - ICU bed reservation.
 - Fasting time
 - Ordering Pre-medication.



Patient Name:
Age: Sex:
Hospital No. :
Date:

1- Pre-Operative Assessment Note

Patient seen in Pre-operative Anesthesia Clinic? ☐ YES ☐ NO

A- History

Previous Anesthesia:	Cardiovascular:	Weight kg	Height: cm
Complications?		Age:	
Airway Difficulty?		Past Med. Hx.:	
Allergies:	Respiratory:	Fasting Status:	
Medications:		Other:	

B- Physical Examination

Vital Signs:	Cardiovascular:	Other:
B/P		
Pulse	Respiratory:	
Temp		
R/R	Airway: H&N movement:	
Pain	Mouth Opening: Tracheal Shift? Mallampati class: Thyromental Distance: Teeth: Other:	

C- Investigations:

Full Blood Count:	Chest X Ray:	Other:
Electrolytes:		
Arterial Blood Gases:	ECG:	

D- Assessment Outcome:

ASA:	Possible Modalities of Anesthesia	Anesthesia & Pain management Plan discussed with Patient/ Family? ☐ YES ☐ NO	Consent taken yet? ☐ YES ☐ NO
-------------	--	--	---

E- PLAN:

Anesthetist's Name:	Signature :	Date/ Time:
----------------------------	--------------------	--------------------

Informed Consent

The following potential complications of General Anesthesia or Regional Anesthesia have been thoroughly explained to me: - Nausea, vomiting - Headache - Sore throat - Damage of the teeth and eyes - Backache - Paresis – Post Peripheral Nerve Block after regional anesthesia - Aspiration (if not fasting) - Cardiac Arrest - Brain Damage - Nerve damage after Spinal Injection - Nerve damage after Epidural Injection - Bleeding or hematoma at the site of spinal or epidural injection. - Infection at the site of spinal or epidural injection or meningitis. - Venous thrombosis - Chronic pain at the site of spinal or epidural injection. - Respiratory and cardiovascular depression.	لقد تم شرح احتمالية حدوث الاعراض الجانبية التالية من إجراء التخدير بتواضع: - القيء، الغثيان - الصداع - التهاب الحلق - تلف الأسنان والعيون - الآلام في الظهر - مرحلة ما بعد احصار العصب المحيطي – شلل جزئي - الارتجاج - سكتة قلبية - تلف في الدماغ - تلف في الأعصاب بعد: تخدير الحبل الشوكي - تلف في الأعصاب بعد: تخدير فوق الجافية - نزف في مكان التخدير الحبل الشوكي أو فوق الجافية - التهاب في مكان التخدير الحبل الشوكي أو فوق الجافية - تجلط دموي - الم مزمن في مكان التخدير الحبل الشوكي أو فوق الجافية - تتبع في عمل الجهاز التنفسي أو القلبي الوعائي
I understand that the anesthetic plan chosen by my anesthesia team is the most appropriate in my case and that my anesthetic management will be provided by adequately privileged doctor. I further understand that whenever the type of anesthesia needs to be altered from what has already been discussed, it is for the purpose of my safety and wellbeing I authorize the anesthesia team to transfuse blood and blood products based on my clinical needs during the treatment procedure	لقد فهمت بأن التخدير سيتم إعطاؤه من قبل طبيب مؤهل من فريق التخدير كما سيقوم طبيب تخدير باختيار طريقة التخدير الأكثر امان وأكثر فاعلية. كما أنني أعلم أنه إذا استدعت الظروف قد يتم تغيير طريقة التخدير عما تمت مناقشته مسبقاً، فإنه سيكون الهدف من ذلك سلامتي وصحتي. أخول فريق التخدير بنقل الدم ومشتقاته إذا استدعت الحاجة إلى ذلك أثناء الإجراء الطبي.
I understand that although favorable results can be expected, they cannot be guaranteed. There is no guarantee against poor results or complications of anesthesia.	أعلم أنه بالرغم من توقع نتائج مرجوة، فإنه لا يمكن ضمانها. حيث أنه لا ضمان من عدم وقوع مضاعفات أو نتائج سلبية من التخدير.
I confirm that I have read and fully understand the above.	أؤكد أنني قرأت وفهمت كل ما ورد أعلاه.
Anesthetist Name & Signature Date: Time:	اسم وتوقيع طبيب التخدير التاريخ الوقت.....
Patient's/ Guardian's Signature الوصي القانوني / المريض	
Patient Guardian's Relationship to Patient and his/her name صلة الوصي القانوني بالمريض واسمه يرجى التوضيح	

 مستشفى الجامعة الأردنية Jordan University Hospital CLN 0232 Patient ID Label Patient Name : File no.:	
WRITTEN CONSENT FOR ANAESTHESIA I (Name) confirm that Dr. has explained to me that I will require: Anesthesia under the care of anesthesia team Headed by..... For (operation/procedure) To be performed on (date of operation/ procedure)	الموافقة الخطية على التخدير أقر أنا (الاسم) العنوان أن الطبيب قد شرح لي أنني بحاجة إلى: التخدير تحت إشراف فريق التخدير برئاسة الدكتور من أجل (العملية/ الإجراء) وذلك بتاريخ:
The Anesthesia proposed for this operation / procedure is: (tick as appropriate). ☐ General Anesthesia ☐ Combined Spinal Epidural (CSE) ☐ Spinal Anesthesia ☐ Epidural Anesthesia ☐ Peripheral Nerve Block (Plexus) ☐ Sedation ☐ Local Anesthesia I understand that Anesthesia through the plan might change according to the condition of operation and patient's safety.	الرجاء وضع علامة على نوع التخدير المقترح للعملية / الإجراء. ☐ التخدير العام ☐ تخدير فوق الجافية المزودة ☐ تخدير الحبل الشوكي ☐ تخدير فوق الجافية ☐ تخدير الأعصاب الطرفية ☐ التشنج الدوائي والتتويم العميق ☐ تخدير موضعي علماً بأنه قد يتغير نوع التخدير المقترح . ويتحول إلى نوع آخر حسب متطلبات العملية وسلامة المريض.
The Anesthetist has fully explained to me the technique and the associated risks (both during the administration of the anesthesia and during the recovery period), benefits and possible alternatives. I have been given an opportunity to ask questions and all of my questions have been answered fully.	لقد شرح لي طبيب التخدير شرحاً وافياً عن كيفية التخدير المشار إليه والمخاطر المتعلقة به أثناء إعطاء التخدير وأثناء الإفاقة من التخدير وفوائد التخدير والبدائل الممكنة. ولقد تم إعطائي الفرصة لطرح الأسئلة وتمت الإجابة على أسئلتني بشكل وافٍ.
The following specific high risks were explained in my case :	لقد شرح لي طبيب التخدير المخاطر المحددة المتعلقة بحالتي التالية :



Prevention of aspiration ASA **Fasting** Guidelines

Clear fluid	2 hours	Water , Fruit juice without pulp,
Milk		
Human	4 hours	
Infant formula	6 hours	
Light Foods	6 hours	Fruits , juice with pulp, Vegetables
Heavy foods	8 hours	Fatty meals , meats

Premedication

- Anxiolysis
- Prevention of aspiration
- DVT / PE prophylaxis
- Antibiotics prophylaxis

Anxiolysis

- ❑ Visit and interview and establishing good rapport with patient.
- ❑ Medications:
 - Benzodiazepines:
 - Diazepam,
 - Midazolam: I.V. (shorter acting than Diazepam) (oral liquid form for children)
 - Opioids:
 - Morphine
 - Pethidine

Perioperative Aspiration

- Risk factors:
 - Fasting state: incomplete
 - Surgical condition:
 - Intestinal obstruction
 - Severe Pain
 - Medications: opioid use
 - Patient related
 - Obesity
 - Gastro-esophageal Reflux disease (GERD)
 - Hiatus Hernia
 - Pregnancy

Drugs used to decrease incidence of aspiration

□ H2 Blockers

- Classes include Cimetidine, Ranitidine (Zantac), Famotidine.
- They block histamine receptor ability to induce acid secretion by proton pump.
 - they consequently reduce gastric fluid volume and acidity

□ Antacids

- Given ½ an hour before induction : 30 ml of sodium citrate
 - Reduce gastric acidity only

□ PPI

- Omeprazole, lansoprazole, and esomeprazole .
 - ✓ Binds to H⁺ / K⁺ pump on parietal cell.
 - ✓ Given 40 mg IV 30 min before surgery .
 - ✓ Reduce both volume and acidity

□ Metoclopramide

- Act on dopamine receptors
- increase gastric motility & lower esophageal sphincter tone
- Reduce gastric fluid volume only

Risk factors for intraoperative DVT

- History of DVT
- **Hypercoagulable states:** Antithrombin III deficiency, Protein C deficiency, Protein S deficiency, Plasminogen activator deficiency.
- Prolonged preoperative immobility.
- Oral contraceptives, pregnancy, post-partum state.
- Long bone fractures.
- Pelvic and lower extremity surgeries.
- Cancer
- Heart failure
- Obesity
- Smoking
- Prolonged surgery
- *Etc.*

THROMBOSIS RISK FACTOR ASSESSMENT

CHOOSE ALL THAT APPLY

EACH RISK FACTOR REPRESENTS 1 POINT

- ☐ Age 41-60 years
- ☐ Minor surgery planned
- ☐ History of prior major surgery (< 1 month)
- ☐ Varicose veins
- ☐ History of inflammatory bowel disease
- ☐ Swollen legs (current)
- ☐ Obesity (BMI > 25)
- ☐ Acute myocardial infarction
- ☐ Congestive heart failure (<1 month)
- ☐ Sepsis (<1 month)
- ☐ Serious lung disease including pneumonia (<1 month)
- ☐ Abnormal pulmonary function (COPD)
- ☐ Medical patient currently at bed rest
- ☐ Other risk factors

EACH RISK FACTOR REPRESENTS 2 POINTS

- ☐ Age 60-74 years
- ☐ Arthroscopic surgery
- ☐ Malignancy (present or previous)
- ☐ Major surgery (> 45 minutes)
- ☐ Laparoscopic surgery (> 45 minutes)
- ☐ Patient confined to bed (> 72 hours)
- ☐ Immobilizing plaster cast (< 1 month)
- ☐ Central venous access

EACH RISK FACTOR REPRESENTS 3 POINTS

- ☐ Age over 75 years
- ☐ History of DVT/PE
- ☐ Family history of thrombosis*
- ☐ Positive Factor V Leiden
- ☐ Positive Prothrombin 20210A
- ☐ Elevated serum homocysteine
- ☐ Positive lupus anticoagulant
- ☐ Elevated anticardiolipin antibodies
- ☐ Heparin-induced thrombocytopenia (HIT)
- ☐ Other congenital or acquired thrombophilia

If yes:

Type: _____

* most frequently missed risk factor

EACH RISK FACTOR REPRESENTS 5 POINTS

- ☐ Elective major lower extremity arthroplasty
- ☐ Hip, pelvis or leg fracture (< 1 month)
- ☐ Stroke (< 1 month)
- ☐ Multiple trauma (< 1 month)
- ☐ Acute spinal cord injury (paralysis) (< 1 month)

FOR WOMEN ONLY (EACH REPRESENTS 1 POINT)

- ☐ Oral contraceptives or hormone replacement therapy
- ☐ Pregnancy or postpartum (< 1 month)
- ☐ History of unexplained stillborn infant, recurrent spontaneous abortion ≥ 3 , premature birth with toxemia or growth-restricted infant

TOTAL RISK FACTOR SCORE

2005 Caprini Risk Assessment Model
Reprinted with permission from
Joseph A. Caprini, MD

■ Interpretation:

Total Risk Factor Score	Risk Level	DVT Incidence	Prophylaxis Regimen*
0 - 1	Low	<10%	No specific measures; early ambulation
2	Moderate	10 - 20%	ES, IPC, LDUH (5000U BID), or LMWH (<3400 U)
3 - 4	High	20 - 40%	IPC, LDUH (5000U TID), or LMWH (>3400U)
≥5	Highest	40 - 80% with 1 - 5% mortality	Pharmacological: LDUH, LMWH (>3400 U), Warfarin, or FXa I alone or in combination with ES or IP

*Where: ■ ES – elastic stockings ■ IPC – intermittent pneumatic compression ■ LDUH – low dose unfractionated Heparin ■ LMWH – low molecular weight Heparin ■ FXa I – Factor X inhibitor

Disclaimer: This tool should NOT be considered as a substitute for any professional medical service, NOR as a substitute for clinical judgement.

Antibiotics Prophylaxis

- ❑ Prevention of surgical site infection.
- ❑ Prevention of infection in immunocompromised patients.
- ❑ Best time for administration is 60 minutes before the surgery (better tissue concentration).
 - Two exceptions for this rule
 - Vancomycin: before 2 hours
 - Use of Tourniquet : prior to inflation
- ❑ Re-dosing in very long surgeries.



Prevention of surgical site infection (SSI)

- Choice of antibiotic is surgery dependent, patient dependent (kidney and liver function), and Hospital dependent (specific types of antibiotics depends on presence of local strains of bacteria resistant/susceptible to common antibiotics).

Table 1. Antibiotic Prophylaxis to Prevent Surgical Site Infections

<i>Surgery</i>	<i>Common pathogens</i>	<i>Recommended antimicrobials*</i>
Cardiothoracic	<i>Staphylococcus aureus</i> , coagulase-negative staphylococci	Cefazolin, cefuroxime sodium (Zinacef), or vancomycin
Gastrointestinal	Enteric gram-negative bacteria, anaerobes, enterococci	Cefoxitin (Mefoxin), cefotetan (Cefotan), ampicillin/sulbactam (Unasyn), or cefazolin plus metronidazole
Gynecologic (vaginal, abdominal, or laparoscopic hysterectomy)	Enteric gram-negative bacteria, group B streptococci, enterococci, anaerobes	Cefoxitin, cefotetan, cefazolin, or ampicillin/sulbactam
Orthopedic	<i>S. aureus</i> , coagulase-negative staphylococci	Cefazolin, cefuroxime sodium, or vancomycin
Vascular	<i>S. aureus</i> , coagulase-negative staphylococci, enteric gram-negative bacilli	Cefazolin or vancomycin

*—Antibiotics are given intravenously within one hour before surgery, except for vancomycin or fluoroquinolones (infusion should start one to two hours before incision). Some authors recommend weight-based dosing of cephalosporins and vancomycin: cephalosporins, 1 g for patients weighing < 176 lb (80 kg) and 2 g for patients weighing ≥ 176 lb; vancomycin, 1 g or 15 mg per kg for patients weighing > 165 lb (75 kg) up to a maximum of 1.5 g. Ampicillin/sulbactam should be administered as a standard 3 g dose. Metronidazole can be administered as a 0.5 g to 1.0 g dose. For patients with normal renal function, an additional intraoperative dose of antibiotic can be administered for surgeries lasting more than four hours or if blood loss > 1,500 mL occurs. Redosing intervals should be based on one to two times the half-life of the drug. Vancomycin can be used when methicillin-resistant *S. aureus* or coagulase-negative staphylococci are common causes of postoperative wound infections, for patients allergic to beta-lactam antibiotics, or when clindamycin (Cleocin) is not appropriate therapy. For patients allergic to penicillins and cephalosporins, clindamycin with ciprofloxacin (Cipro), levofloxacin (Levaquin), or aztreonam (Azactam) is a reasonable alternative.

Information from references 8 and 9.

Thank You!

